

Henry Ford Health System

Henry Ford Health System Scholarly Commons

Clinical Research

Medical Education Research Forum 2019

5-2019

Intravenous and Oral Tranexamic Acid are Equivalent at Reducing Blood Loss in Thoracolumbar Spinal Fusion: A Prospective Randomized Trial Phase 2

Charles C. Yu

Henry Ford Health System, CYu1@hfhs.org

Omar M Kadri

Henry Ford Health System, OKADRI1@hfhs.org

Allen Kadado

Henry Ford Health System, akadado1@hfhs.org

Jacob Pawloski

Henry Ford Health System, JPAWLOS1@hfhs.org

Morenikeji Buraimoh

Henry Ford Health System

See next page for additional authors

Follow this and additional works at: <https://scholarlycommons.henryford.com/merf2019clinres>

Recommended Citation

Yu, Charles C.; Kadri, Omar M; Kadado, Allen; Pawloski, Jacob; Buraimoh, Morenikeji; Bartol, Stephen; and Graziano, Gregory, "Intravenous and Oral Tranexamic Acid are Equivalent at Reducing Blood Loss in Thoracolumbar Spinal Fusion: A Prospective Randomized Trial Phase 2" (2019). *Clinical Research*. 41. <https://scholarlycommons.henryford.com/merf2019clinres/41>

This Poster is brought to you for free and open access by the Medical Education Research Forum 2019 at Henry Ford Health System Scholarly Commons. It has been accepted for inclusion in Clinical Research by an authorized administrator of Henry Ford Health System Scholarly Commons.

Authors

Charles C. Yu, Omar M Kadri, Allen Kadado, Jacob Pawloski, Morenikeji Buraimoh, Stephen Bartol, and Gregory Graziano

Intravenous and oral tranexamic acid are equivalent at reducing blood loss in thoracolumbar spinal fusion: a prospective randomized trial phase 2

Charles C. Yu, MD, Omar Kadri, MD, Allen Kadado, MD, Jacob Pawloski, BS, Morenikeji Buraimoh, MD, Stephen Bartol, MD, Gregory Graziano, MD.

Henry Ford Health System



May 3rd, 2019



all for you


Background

- Tranexamic acid (TXA) is a synthetic antifibrinolytic that prevents fibrin breakdown by binding to plasminogen, thus stabilizing the fibrin clot.¹
- Intraoperative use of TXA has proven to be effective in reducing transfusion requirements after spinal fusion.^{2,3}



Background

- TXA can be administered orally, topically, or IV.
- IV TXA costs \$47-\$108 per dose depending on formulation.¹⁰
- Oral TXA is both cheaper and easier to administer than IV.
- There are currently no prospective studies investigating equivalence between IV and PO TXA in spinal fusion surgery.



Methods

- PO TXA: 1950mg (3 tablets) 2 hrs before incision
- IV TXA: 1g (in 100mL NS) bolus before incision and 1g before closure
- 2 surgeons: G.G. performed majority of surgeries (95%)
- Revision cases: 63% of total
- Exclusion criteria: known allergy to TXA, history of renal failure or kidney transplant, history of arterial thromboembolic event (eg. myocardial infarction, stroke) within the past year, placement of an arterial stent within the past year, a history of thromboembolic event, coagulopathy, or refusal to receive blood products.

A randomized, controlled trial

Enrollment

Assessed for eligibility
(n = 200)

Excluded (n = 29)
Not meeting inclusion criteria
(n = 16)
Refused to participate
(n = 6)
Other reasons (n = 7)

Randomized (n = 171)

Allocation

Allocated to IV TXA
(n = 91)
Received allocated
intervention (n = 91)

Allocated to PO TXA
(n = 80)
Received allocated
intervention (n = 80)

Follow up

Lost to follow up
(n = 0)
Discontinued
intervention (n = 0)

Lost to follow up
(n = 0)
Discontinued
intervention (n = 0)

Analysis

Analyzed (n = 91)

Analyzed (n = 80)

Table 1. Patient characteristics

	IV	PO	P*
Number of patients	91	80	
Mean age (yr)	64 (13)	60 (13)	0.02
Sex (M/F)	40/51	34/46	0.85 ^a
Weight (kg)	85 (20)	92 (21)	0.02
Height (m)	1.69 (0.10)	1.69 (0.11)	0.7
Body mass index	29.4 (6.0)	32.1 (5.8)	0.003
Estimated blood volume (mL)	4903 (941)	5107 (1035)	0.18
ASA	2.7 (0.5)	2.8 (0.5)	0.67
Class I (n)	0	1	
Class II (n)	24	19	
Class III (n)	64	60	
Class IV (n)	3	0	
Number of fused levels	3.8 (3.0)	3.6 (2.9)	0.82
1-2 levels	42	42	
3-5 levels	27	21	
>5 levels	22	17	
Number (%) of revisions	59 (65%)	49 (61%)	0.63 ^a
Number (%) of interbody fusions	9 (10%)	19 (24%)	0.01 ^a
Number (%) of osteotomies	13 (14%)	12 (15%)	0.90 ^a
Number (%) of anticoagulant use	11 (12%)	2 (5%)	0.19 ^a
Preoperative Hgb (g/dL)	13.2 (1.5)	13.3 (1.7)	0.54
Preoperative Hct (%)	39.4 (4.5)	39.9 (5.1)	0.48
Preoperative platelet count (x10 ³ /mm ³)	208 (58)	254 (72)	<0.001
Preoperative INR	1.09 (0.14)	1.06 (0.19)	0.39
Preoperative creatinine (mg/dL)	0.99 (0.82)	0.94 (0.87)	0.7

Mean (SD)

*Statistical analysis for comparison between groups: P < 0.05, statistical significance

^ax² test or Fisher exact test.



all for you

Table 1. Patient characteristics

	IV	PO	P*
Number of patients	91	80	
Mean age (yr)	64 (13)	60 (13)	0.02
Sex (M/F)	40/51	34/46	0.85 ^a
Weight (kg)	85 (20)	92 (21)	0.02
Height (m)	1.69 (0.10)	1.69 (0.11)	0.7
Body mass index	29.4 (6.0)	32.1 (5.8)	0.003
Estimated blood volume (mL)	4903 (941)	5107 (1035)	0.18
ASA	2.7 (0.5)	2.8 (0.5)	0.67
Class I (n)	0	1	
Class II (n)	24	19	
Class III (n)	64	60	
Class IV (n)	3	0	
Number of fused levels	3.8 (3.0)	3.6 (2.9)	0.82
1-2 levels	42	42	
3-5 levels	27	21	
>5 levels	22	17	
Number (%) of revisions	59 (65%)	49 (61%)	0.63 ^a
Number (%) of interbody fusions	9 (10%)	19 (24%)	0.01 ^a
Number (%) of osteotomies	13 (14%)	12 (15%)	0.90 ^a
Number (%) of anticoagulant use	11 (12%)	2 (5%)	0.19 ^a
Preoperative Hgb (g/dL)	13.2 (1.5)	13.3 (1.7)	0.54
Preoperative Hct (%)	39.4 (4.5)	39.9 (5.1)	0.48
Preoperative platelet count (x10 ³ /mm ³)	208 (58)	254 (72)	<0.001
Preoperative INR	1.09 (0.14)	1.06 (0.19)	0.39
Preoperative creatinine (mg/dL)	0.99 (0.82)	0.94 (0.87)	0.7
Mean (SD)			

*Statistical analysis for comparison between groups: P < 0.05, statistical significance

^ax² test or Fisher exact test.



all for you

Results

Table 2. Intraoperative measures

	IV (N=91)	PO (N=80)	P*
Anesthesia time (min)	360 (109)	357 (109)	0.86
Surgical time (min)	261 (106)	265 (106)	0.79
Estimated blood loss (mL)	350 (50, 3000)	325 (40, 2300)	0.83 ^b
Intravenous fluid (mL)	3198 (1589)	3387 (1610)	0.44
Number (%) receiving cell saver	29 (32%)	29 (36%)	0.55 ^a
Number (%) receiving blood transfusion	8 (9%)	3 (4%)	0.18 ^a

Mean (SD)

EBL presented as median (min, max) due to nonparametric distribution.

*Statistical analysis for comparison between groups: $P < 0.05$, statistical significance.

^a χ^2 test or Fisher exact test.

^bWilcoxon sum run test.

Results

Table 3. Outcome measurements

	IV (N=91)	PO (N=80)	P*
Hgb drop (g/dL)	3.48 (1.85)	3.19 (1.54)	0.004 ^c
Calculated blood loss (mL)	1274 (688)	1206 (622)	0.001 ^c
Hct drop (%)	10.3 (5.6)	9.7 (5.7)	0.003 ^c
Drain output (mL)	637 (459)	641 (491)	0.95
Rate of postop transfusion	23%	13%	0.07 ^a
Rate of DVT/PE	2%	3%	0.90 ^a
Rate of infections	3%	4%	0.41 ^a
Length of hospital stay (days)	4.5 (3.2)	3.8 (2.7)	0.11

Mean (SD)

*Statistical analysis for comparison between groups: $P < 0.05$, statistical significance.

^a χ^2 test or Fisher exact test.

^cWelch two one-sided test: $P < 0.05$ demonstrates equivalence between treatments.

Results

Table 3. Outcome measurements

	IV (N=91)	PO (N=80)	P*
Hgb drop (g/dL)	3.48 (1.85)	3.19 (1.54)	0.004 ^c
Calculated blood loss (mL)	1274 (688)	1206 (622)	0.001 ^c
Hct drop (%)	10.3 (5.6)	9.7 (5.7)	0.003 ^c
Drain output (mL)	637 (459)	641 (491)	0.95
Rate of postop transfusion	23%	13%	0.07 ^a
Rate of DVT/PE	2%	3%	0.90 ^a
Rate of infections	3%	4%	0.41 ^a
Length of hospital stay (days)	4.5 (3.2)	3.8 (2.7)	0.11

Mean (SD)

*Statistical analysis for comparison between groups: P < 0.05, statistical significance.

^ax² test or Fisher exact test.

^cWelch two one-sided test: P < 0.05 demonstrates equivalence between treatments.

Discussion

- PO TXA has been shown to be equivalent to IV in total joints literature
- First study comparing PO and IV TXA in thoracolumbar fusions
- Use of oral TXA in spinal fusion surgery could provide substantial savings to the healthcare system
- At least \$20 million per year of saving

Limitations

- Heterogeneity in patient diagnosis and surgical technique/approach
- Hgb change was based on lowest recorded post-operative hgb, which could be affected by hemodilution
- No placebo group and assumed PO TXA was superior than placebo

Summary

- Oral TXA is equivalent to IV TXA in preventing blood loss and hemoglobin drop in spinal fusion surgery.
- There is no significant difference in drain output, post-op transfusions, DVT/PE, or infections when using IV or PO TXA.
- Oral TXA is a viable, more cost-effective alternative to IV TXA to prevent blood loss in spine surgery.

Future Directions

- Ongoing trial to collect a larger cohort to perform adequately powered sub-group analysis

References

- Mannucci PM: **Hemostatic drugs.** *N Engl J Med* 1998, **339**(4):245-253.
- Gill JB, Chin Y, Levin A, Feng D: **The use of antifibrinolytic agents in spine surgery. A meta-analysis.** *J Bone Joint Surg Am* 2008, **90**(11):2399-2407.
- Li ZJ, Fu X, Xing D, Zhang HF, Zang JC, Ma XL: **Is tranexamic acid effective and safe in spinal surgery? A meta-analysis of randomized controlled trials.** *Eur Spine J* 2013, **22**(9):1950-1957.
- Fillingham YA, Kayupov E, Plummer DR, Moric M, Gerlinger TL, Della Valle CJ:
- **The James A. Rand Young Investigator's Award: A Randomized Controlled Trial of Oral and Intravenous Tranexamic Acid in Total Knee Arthroplasty: The Same Efficacy at Lower Cost?** *J Arthroplasty* 2016, **31**(9 Suppl):26-30.

Thank you



all for you