

Henry Ford Health

## Henry Ford Health Scholarly Commons

---

Hypertension and Vascular Research Articles

Hypertension and Vascular Research

---

8-17-2022

### **Nitric oxide-inhibited chloride transport in cortical thick ascending limbs is reversed by 8-iso-prostaglandin-F2 $\alpha$**

Pablo D. Cabral

Guillermo B. Silva

Sandra T. Baigorria

Luis I. Juncos

Ebenezer I. O Ajayi

*See next page for additional authors*

Follow this and additional works at: [https://scholarlycommons.henryford.com/hypertension\\_articles](https://scholarlycommons.henryford.com/hypertension_articles)

---

---

**Authors**

Pablo D. Cabral, Guillermo B. Silva, Sandra T. Baigorria, Luis I. Juncos, Ebenezer I. O Ajayi, and Néstor H. García

# Nitric oxide-inhibited chloride transport in cortical thick ascending limbs is reversed by 8-iso-prostaglandin-F2 $\alpha$

Pablo D. Cabral<sup>1</sup>, Guillermo B. Silva<sup>2,3</sup>, Sandra T. Baigorria<sup>2</sup>, Luis I. Juncos<sup>2</sup>, Ebenezer I. O. Ajayi<sup>4,5</sup>, Néstor H. García<sup>2</sup>

<sup>1</sup>Hypertension and Vascular Research Division, Department of Internal Medicine, Henry Ford Hospital, Detroit, MI, USA

<sup>2</sup>Department of Renal Physiology, J. Robert Cade Foundation, Córdoba, Argentina

<sup>3</sup>Department of Renal Physiology and Hypertension, Mons. Carlos V. Cruvellier Foundation, San Juan, Argentina

<sup>4</sup>DC&ONID, Biochemistry Department, Osun State University, Osogbo, Nigeria

<sup>5</sup>IMMF-INIMEC-UNC, CONICET, Córdoba, Argentina

**Background:** Sodium chloride (NaCl) reabsorption in the cortical thick ascending limb (cTAL) is regulated by opposing effects. Nitric oxide (NO) inhibits NaCl reabsorption while 8-iso-prostaglandin-F2 $\alpha$  (8-iso-PGF2 $\alpha$ ) stimulates it. Their interaction has not been evaluated in the cTAL. Because 8-iso-PGF2 $\alpha$  has considerable stability while NO is a free radical with a short half-life, we hypothesized that, in the cTAL, the inhibition of NaCl absorption will be reversed by 8-iso-PGF2 $\alpha$ .

**Methods:** Chloride absorption ( $J_{Cl}$ ) was measured in isolated perfused cTALs and whether the activation of protein kinase A (PKA) is required for this interaction. Since cyclic adenosine monophosphate (cAMP) is a major messenger for the 8-iso-PGF2 $\alpha$  signaling cascade, and NO inhibits  $J_{Cl}$  by decreasing cAMP bioavailability, we measured 8-iso-PGF2 $\alpha$ -stimulated cAMP in the presence of sodium nitroprusside (SNP).

**Results:** The NO donor, SNP ( $10^{-6}$  M), decreased  $J_{Cl}$  by 41%, while luminal 8-iso-PGF2 $\alpha$  (100  $\mu$ M) increased  $J_{Cl}$  to  $315 \pm 46$  pmol/min/mm ( $p < 0.003$ ), reversing the effects of the NO donor. SNP inhibited  $J_{Cl}$ , 8-iso-PGF2 $\alpha$  failed to increase  $J_{Cl}$  in the presence of H89. Basal cAMP was  $56 \pm 13$  fmol/min/mm, in the presence of SNP  $57 \pm 6$  fmol/min/mm, and 8-iso-PGF2 $\alpha$  increased it to  $92 \pm 2$  fmol/min/mm ( $p < 0.04$ ).

**Conclusion:** We concluded that 1) NO-induced inhibition of  $J_{Cl}$  in the cTAL can be reversed by 8-iso-PGF2 $\alpha$ , 2) 8-iso-PGF2 $\alpha$  and NO interaction requires PKA to control  $J_{Cl}$ , and 3) in the presence of NO, 8-iso-PGF2 $\alpha$  continues to stimulate  $J_{Cl}$  because NO cannot reverse 8-iso-PGF2 $\alpha$ -stimulated cAMP level.

**Keywords:** Hypertension, Isoprostanes, Nitric oxide, Oxidative stress

## Introduction

The nephron segment, cortical thick ascending limb (cTAL)

regulates sodium chloride (NaCl) reabsorption as one of the most involved segments in NaCl retention states. In normal conditions, absorption of NaCl is balanced with

**Received:** October 22, 2021; **Revised:** March 14, 2022; **Accepted:** April 11, 2022

**Correspondence:** Néstor H. García

Department of Renal Physiology, J. Robert Cade Foundation, Pedro de Oñate 253, Córdoba 5003, Argentina. E-mail: [garcia.nestor@conicet.gov.ar](mailto:garcia.nestor@conicet.gov.ar)  
ORCID: <https://orcid.org/0000-0002-9057-9030>

This paper is dedicated to the memory of Guillermo B. Silva.

Copyright © 2022 by The Korean Society of Nephrology

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial and No Derivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>) which permits unrestricted non-commercial use, distribution of the material without any modifications, and reproduction in any medium, provided the original works properly cited.

its excretion, maintaining Na balance. This homeostasis is the final result of the action of molecules increasing and decreasing transport. However, in pathologic conditions, such as congestive heart failure, chronic kidney disease, hypertension, cirrhosis, and other diseases, Na retention is greater than its excretion, leading to a positive Na imbalance. This Na overload occurs because Na retention factors overcome the actions of molecules that increase its excretion, resulting in resistance to natriuretic effects. Understanding the interaction of the autacoids at the different nephron segments, especially at the cTAL, would help to elucidate such deleterious conditions and find strategies to manage them.

Previously, we reported that 8-iso-prostaglandin-F<sub>2</sub> $\alpha$  (8-iso-PGF<sub>2</sub> $\alpha$ ) increases Na<sup>+</sup> and Cl<sup>-</sup> reabsorption at the cTAL [1]. This effect, *in vivo*, would increase Na balance, inducing a blood pressure elevation that could be displayed as hypertension, congestive heart failure, cirrhosis, chronic kidney disease, or other edematous states. On the other hand, nitric oxide (NO) inhibits Na<sup>+</sup> reabsorption at the cTAL, and this effect, *in vivo*, would induce diuresis and natriuresis, a mechanism that would decrease the described Na gain [2,3]. Despite the clear understanding of the importance of these effects, the interaction of NO and 8-iso-PGF<sub>2</sub> $\alpha$  in the cTAL has not been evaluated to elucidate their outcomes.

Na<sup>+</sup> regulation has been previously reported at several nephron segments mediated by agents such as NO and angiotensin II in the cTAL [4,5], NO and arginine vasopressin at the cortical collecting duct level [6], and other segments [7]. The final effect of the interaction between NO and 8-iso-PGF<sub>2</sub> $\alpha$  is based on the intracellular activation of different regulators. Actually, whereas NO inhibits NaCl reabsorption by increasing intracellular cyclic guanosine monophosphate (cGMP) and decreasing cyclic adenosine monophosphate (cAMP) levels [7], 8-iso-PGF<sub>2</sub> $\alpha$  increases cAMP. The final effect will be elicited when one cyclic nucleotide preponderantly represses the other. When attempting to understand the crosstalk regulation of cyclic nucleotide concentrations, one must consider that such regulation can occur at the level of cyclic nucleotide synthesis, phosphodiesterase-mediated degradation, or even cellular efflux. Thus, we hypothesized that, in cTAL, persistent effects of 8-iso-PGF<sub>2</sub> $\alpha$  on Na reabsorption prevail

over the shorter-acting NO effects, and protein kinase A (PKA) activation is required for such interaction.

## Methods

### Animals

Male New Zealand white rabbits (700–1,200 g) were housed for 3 to 7 days before the study for acclimatization. The experiments were approved by the Institutional Animal Care and Use Committee of the J. Robert Cade Foundation (No. FJRC #2009) in Córdoba, Argentina. All animals were housed and handled according to the Public Health Service Policy on Humane Care and Use of Laboratory Animals.

### Preparation and dissection of the cortical thick ascending limb of Henle

Animals were anesthetized with sodium thiopental (30 mg/kg, intravenously) and xylazine (20 mg/kg, intramuscularly) plus ketamine (50 mg/kg, intramuscularly). The kidney was exposed and bathed in ice-cold saline and then excised and cut into coronal slices along the longitudinal axis. The slices were placed in oxygenated physiological saline in a temperature-regulated chamber at 4°C. The cTAL of Henle was dissected from the medullary ray under a stereomicroscope as previously described [6,8].

### Perfusion of thick ascending limbs

cTALs in a temperature-regulated chamber were perfused between concentric glass pipettes at 37°C as described [6]. The basolateral bath was exchanged at a rate of 0.5 mL/min. A total of five to eight tubules were analyzed during each protocol. Time-control experiments were conducted to ensure the stability of tubular transport. Chloride absorption ( $J_{Cl}$ ) was measured as previously described [6,9]. cTALs were perfused and equilibrated for 20 minutes, and basal Cl absorption was calculated two to three times. Tested compounds, including 8-iso-PGF<sub>2</sub> $\alpha$ , were either added to or removed from the lumen or bath as indicated. After a 20-minute equilibration period, tubular fluid was again collected 2 or 3 times.

## Chloride concentrations in the perfusate and collected fluid

Chloride concentrations were measured by microfluorometry [10]. Because water is not reabsorbed by the thick ascending limb, J<sub>Cl</sub> was calculated by the following equation:  $J_{Cl} = CR \times (C_{0Cl} - C_{1Cl})$ , where CR = collection rate normalized/tubule length, C<sub>0Cl</sub> = chloride concentration in the perfusion solution, and C<sub>1Cl</sub> = chloride concentration in the collected fluid.

## Cyclic adenosine monophosphate

cAMP was measured as previously described [1]. Briefly, tubules were isolated and incubated in 95  $\mu$ L perfusion solution containing 1-mM 3-isobutyl-1-methylxanthine at 37°C for 10 minutes before adding 5- $\mu$ L medium containing inhibitors or hormones. The reaction was stopped 30 minutes later with methanol (100  $\mu$ L), and cAMP and cGMP levels were measured by an enzyme immunoassay (EIA; Cayman Chemical, Ann Arbor, MI, USA). The samples were centrifuged, the supernatant was transferred to another tube and dried in a Savant centrifugal vacuum concentrator, and the pellet was reconstituted in 110- $\mu$ L sodium acetate buffer. The cAMP standard was treated the same way. The results were expressed as fmol/min/mm tubule length.

## Statistics


Data were analyzed by paired and unpaired Student t tests and one-way analysis of variance, and results with  $p < 0.05$  were significant. Results were expressed as mean  $\pm$  standard error of the mean.

## Results


Since the 8-iso-PGF2 $\alpha$ /NO interaction in cTAL is uncertain, we chose a maximum NO bioavailability model. As described, the micro-isolated rabbit cTALs were transferred to a temperature-regulated chamber and perfused between concentric glass pipettes at 37°C [6] with the NO donor sodium nitroprusside (SNP) in a concentration that decreases transport by 40% as endogenous NO is inhibited in this nephron segment [11]. Before adding SNP to the

bath, J<sub>Cl</sub> was  $333 \pm 35$  pmol/min/mm. SNP (1 mM) in the bath reduced J<sub>Cl</sub> to  $195 \pm 26$  pmol/min/mm ( $p = 0.01$  vs. basal;  $n = 5$ ) (Fig. 1), a 41% decrease, and remained inhibited for the rest of the experiment ( $196 \pm 22$  pmol/min/mm). *In vivo*, this reduction in J<sub>Cl</sub> should increase diuresis. In time-control experiments, J<sub>Cl</sub> remained constant throughout the experimental period.

Next, we tested whether 8-iso-PGF2 $\alpha$  stimulates J<sub>Cl</sub> in the presence of SNP. Basal J<sub>Cl</sub> was  $330 \pm 18$  pmol/min/mm; when SNP (1 mM) was added to the bath, J<sub>Cl</sub> decreased to  $191 \pm 19$  pmol/min/mm. After adding 8-iso-PGF2 $\alpha$  (100  $\mu$ M) to the lumen, J<sub>Cl</sub> increased to  $346 \pm 39$  pmol/min/mm ( $p < 0.02$  vs. NO;  $n = 5$ ) (Fig. 2). We tested this concentration since those lower did not affect J<sub>Cl</sub> [1]. Next, we evaluated the effect of 8-iso-PGF2 $\alpha$  present in the basolateral side. Basal J<sub>Cl</sub> was  $235 \pm 38$  pmol/min/mm, which decreased in the presence of SNP (1 mM) to  $139.4 \pm 27$  pmol/min/mm and increased to  $297 \pm 29$  pmol/min/mm in the presence of 8-iso-PGF2 $\alpha$  (100  $\mu$ M;  $p < 0.02$  vs. NO;  $n = 5$ ) (Fig. 3). These data indicate that the maximal NO effect on cTAL is insufficient to prevent the 8-iso-PGF2 $\alpha$ -stimulated NaCl reabsorption.




**Figure 1. SNP inhibits J<sub>Cl</sub>; nitric oxide decreases J<sub>Cl</sub> in the cTAL.** Basal J<sub>Cl</sub> was  $333.5 \pm 35.2$  pmol/min/mm and the addition of SNP (1 mM) decreased it to  $195.9 \pm 26.1$  pmol/min/mm (\* $p = 0.01$  vs. basal;  $n = 5$ , one-way analysis of variance [Holm-Sidak test]) and remained decreased during the experimental time ( $196.5 \pm 22.3$  pmol/min/mm). cTAL, cortical thick ascending limb; J<sub>Cl</sub>, chloride absorption; SNP, sodium nitroprusside.



**Figure 2. SNP-induced inhibition of  $J_{Cl}$  was prevented by 8-iso-PGF $2\alpha$  on the luminal side.**

\* $p < 0.003$  vs. basal;  $n = 5$ , one-way analysis of variance (Holm-Sidak test).

$J_{Cl}$ , chloride absorption; SNP, sodium nitroprusside; 8-iso-PGF $2\alpha$ , 8-iso-prostaglandin-F $2\alpha$ .




**Figure 3. SNP-induced inhibition of  $J_{Cl}$  was prevented by 8-iso-PGF $2\alpha$  on the basolateral side.**

\* $p < 0.03$  vs. control period;  $n = 4$ , one-way analysis of variance (Holm-Sidak test).

$J_{Cl}$ , chloride absorption; SNP, sodium nitroprusside; 8-iso-PGF $2\alpha$ , 8-iso-prostaglandin-F $2\alpha$ .

In cTAL, 8-iso-PGF $2\alpha$  stimulates  $J_{Cl}$  via a cAMP-dependent mechanism [1], while NO inhibits chloride reabsorption by stimulation of a cGMP-stimulated phosphodiesterase, which decreases cAMP level [12]. Thus, we evaluated




**Figure 4. Effect of 100  $\mu$ M 8-isoPGF $2\alpha$  on the presence of SNP, a NO donor.**

Basal cAMP was  $56.3 \pm 13.1$  fmol/min/mm, in the presence of the NO donor,  $57.8 \pm 6.1$  fmol/min/mm, and 8-iso-PGF $2\alpha$  increased it to  $92.1 \pm 2.9$  fmol/min/mm (\* $p < 0.001$  vs. control and vs. SNP period;  $n = 10$ , one-way analysis of variance [Tukey honestly significant difference *post hoc* test]).

cAMP, cyclic adenosine monophosphate;  $J_{Cl}$ , chloride absorption; NO, nitric oxide; SNP, sodium nitroprusside; 8-iso-PGF $2\alpha$ , 8-iso-prostaglandin-F $2\alpha$ .

the effect of 8-iso-PGF $2\alpha$  on cAMP in the presence of a NO donor, SNP (1 mM). Basal cAMP was  $56.3 \pm 13.1$  fmol/min/mm, while that in the presence of the NO donor was  $57.8 \pm 6.1$  fmol/min/mm, the addition of 8-iso-PGF $2\alpha$  to which increased cAMP to  $92.1 \pm 2.9$  fmol/min/mm ( $n = 10$ ,  $p < 0.04$ ) (Fig. 4). Thus, 8-iso-PGF $2\alpha$ -stimulated NO-inhibited  $J_{Cl}$  is associated with a 60% increase in cAMP.

In the cTAL, PKA mediates the effect of 8-iso-PGF $2\alpha$  on  $J_{Cl}$ . Thus, we evaluated the effect of 8-iso-PGF $2\alpha$  on SNP-inhibited  $J_{Cl}$  in the presence of H89 (10 mM), a PKA inhibitor. Basal  $J_{Cl}$  was  $249 \pm 13$  pmol/min/mm and decreased to  $173 \pm 14$  pmol/min/mm when SNP (1 mM) was added to the bath. Addition of H89 to the bath did not change  $J_{Cl}$  significantly ( $158 \pm 10$  pmol/min/mm). Under these conditions, 8-iso-PGF $2\alpha$  could not stimulate SNP-inhibited  $J_{Cl}$  ( $147 \pm 15$  pmol/min/mm) (Fig. 5). Time control did not change during the experiment ( $280 \pm 22$  pmol/min/mm vs.  $282 \pm 36$  pmol/min/mm). These data indicate that 8-iso-PGF $2\alpha$  requires PKA activity to reverse the NO-inhibited  $J_{Cl}$ .



**Figure 5. Effect of 8-iso-PGF2 $\alpha$  on NO-induced J<sub>Cl</sub> inhibition in the presence of H89, a PKA inhibitor.**

\*p < 0.03 vs. control period; n = 5, one-way analysis of variance (Holm-Sidak test).

J<sub>Cl</sub>, chloride absorption; NO, nitric oxide; PKA, protein kinase A; 8-iso-PGF2 $\alpha$ , 8-iso-prostaglandin-F2 $\alpha$ .

## Discussion

To our knowledge, this is the first report showing that 8-iso-PGF2 $\alpha$  can override NO's diuretic effects in the cTAL, suggesting that Na<sup>+</sup> retention will prevail over Na<sup>+</sup> excretion in clinical conditions associated with increased 8-iso-PGF2 $\alpha$  level. Although in the kidney, discrete 8-iso-PGF2 $\alpha$  and NO effects have both been shown *in vivo* and *in vitro*, we designed this *in vitro* investigation to identify the local interaction and final effect of these two autacoids on chloride transport at the nephron segment.

Despite the well-known importance of prostaglandin in Na reabsorption, the effects of its derivatives are less clear. We evaluated the effects of 8-iso-PGF2 $\alpha$  binding to the G-protein coupled receptor thromboxane A2 (TXA2) receptor, localized in the thick ascending limb segment [13,14]. This 8-iso-PGF2 $\alpha$  is a modified prostaglandin produced by nonenzymatic oxidation of prostaglandin-F2 $\alpha$  by reactive oxygen species [15,16] and induces vasoconstriction in the kidney and can impact renal blood flow [17]. Our data demonstrate that the NO donor decreases chloride reabsorption in the thick ascending limb of Henle; under these circumstances, 8-iso-PGF2 $\alpha$  is activated by PKA and stimulates Cl<sup>-</sup> reabsorption. *In vivo*, this interaction results in a decrease of natriuresis and sodium increase.

The significance of this interaction resides not only in improving understanding of the normal physiology, but also in the prevention and treatment of many conditions such as hypertension, chronic renal failure, and cirrhosis, where isoprostanes increase and an Na retention state prevails [18].

The compound 8-iso-PGF2 $\alpha$  binds the TXA2 receptor, a peroxidized derivative of prostaglandin-F2 localized in the basolateral and apical membrane of the thick ascending limb segment, which then activates cAMP production [14]. Following the release of 8-iso-PGF2 $\alpha$ , sudden activation of the TXA2 receptor receptor in the thick ascending limb occurs [1], stimulating chloride reabsorption via cAMP. Simultaneously, NO inhibits 8-iso-PGF2 $\alpha$  via cGMP generation. Therefore, the interaction between these two intracellular mechanisms could regulate sodium transport.

In the kidney, 8-iso-PGF2 $\alpha$  induces vasoconstriction, which can modulate renal blood flow [19]. In the thick ascending limb, TXA2 receptor release stimulates PKA and induces an increase in chloride reabsorption across the Na-K-Cl cotransporter 2 (NKCC2) [20]. Increased concentrations of 8-iso-PGF2 $\alpha$  and other isoprostanes have been observed in both the urine and plasma of hypertensive subjects. The enhanced NaCl retention in response to 8-iso-PGF2 $\alpha$  might contribute to the pathogenesis of hypertension [20]. In the distal nephron, NO decreases cAMP, which reduces fluid absorption via PKA [21].

We further investigated whether 8-iso-PGF2 $\alpha$  could reverse NO effects when PKA is inhibited. This experiment should establish whether the interaction between the mechanisms linking 8-iso-PGF2 $\alpha$  and NO precedes or follows activation of PKA. In the presence of PKA inhibition, 8-iso-PGF2 $\alpha$  did not change J<sub>Cl</sub>, indicating that PKA activity is required to reverse the NO-inhibited J<sub>Cl</sub>. Moreover, these results indicate that 8-iso-PGF2 $\alpha$  reverses NO-induced inhibition of J<sub>Cl</sub> via a mechanism involving activation of PKA. Since PKA is mainly activated by cAMP, we studied cAMP and found that NO was not able to reduce the cAMP level induced by 8-iso-PGF2 $\alpha$ .

Nevertheless, the effects of 8-iso-PGF2 $\alpha$  and NO on NaCl excretion have been shown in several models [22]. The decreased urinary Na<sup>+</sup> excretion caused by systemic nitric oxide synthase (NOS) inhibition correlates with higher urinary 8-iso-PGF2 $\alpha$  [23,24]. Also, the superoxide scavenger Tempol blunts these changes, suggesting that inhibition of

NO synthesis enhances endogenous superoxides, which could increase renal 8-iso-PGF2 $\alpha$  level. These studies, however, show correlation but no direct interaction in the transport mechanisms. At any rate, 8-iso-PGF2 $\alpha$  increases Cl<sup>-</sup> by stimulating cAMP synthesis [1].

In contrast, the effect by which NO inhibits NKCC2 activity might involve both an increase in cGMP [12,25,26] and a decrease in cAMP [27]. In our study, the NO donor did not decrease cAMP, whereas cGMP can play an essential role in the NO-inhibited J<sub>Cl</sub>. This effect might involve protein trafficking, limiting the total pool of NKCC2 at the membrane level [28] since 8-iso-PGF2 $\alpha$ 's increase of J<sub>Cl</sub> is mediated by NKCC2 [1].

The mechanism(s) by which cGMP acts to regulate trafficking of the transporter might involve a decrease in insertion of the transporter into the membrane. Although evidence exists supporting the constitutive degradation of surface NKCC2, which is hypothesized to occur by decreasing NKCC2 level due to increasing NKCC2 ubiquitination and proteasomal degradation [28], it appears plausible that NO acts by increasing cGMP, which decreases cAMP [27] to lead to a reduction in exocytic insertion of NKCC2 into the apical membrane, reducing the number of transporters [19]. This process results in lower NKCC2 activity and blunted net NaCl reabsorption.

Our experiments demonstrate that interaction with 8-iso-PGF2 $\alpha$  increases intracellular cAMP; in the presence of H89, 8-iso-PGF2 $\alpha$  does not increase chloride reabsorption. These results show that the interaction occurs after activation of PKA since H89 prevented 8-iso-PGF2 $\alpha$  and J<sub>Cl</sub> inhibited NO and not because 8-iso-PGF2 $\alpha$  blunts cGMP formation. Indeed, we found that NO caused no significant change in 8-iso-PGF2 $\alpha$ -stimulated cAMP level. This indicates that NO-stimulated phosphodiesterase activity has little or no effect on 8-iso-PGF2 $\alpha$ -stimulated cAMP level. This notion might not apply to other cellular compartments, where NO-stimulated phosphodiesterase could be activated separately from 8-iso-PGF2 $\alpha$ -stimulated cAMP. Thus, 8-iso-PGF2 $\alpha$  might interact with NO at the level of cAMP or cGMP.

The pathomechanism whereby 8-iso-PGF2 $\alpha$  prevents NO-inhibited J<sub>Cl</sub> might be observed under several conditions characterized by increased blood pressure, such as hypertension and chronic renal failure. In such conditions, urinary 8-iso-PGF2 $\alpha$  [29] and cAMP [30] are elevated as well as other oxidative stress mediators. On the contrary,

NO production is impaired in part due to limitations on substrate (L-arginine) availability and increased circulating levels of endogenous NO synthase inhibitors, in particular asymmetric dimethylarginine and reduced renal cortex abundance of the neuronal NOS [31]. This effect of NO correlates with urinary cGMP level [32]. This pathological condition is characterized by sodium retention as a consequence of enhanced sympathetic activity [33] and increase synthesis of intrarenal angiotensin II synthesis [34], endothelin-1 [35], and 8-iso-PGF2 $\alpha$ , all of which stimulate sodium reabsorption. In contrast, the antinatriuretic effect of NO is blunted under these conditions, leading to sodium retention, hypertension, fibrosis, and end-organ damage.

The beneficial effect of 8-iso-PGF2 $\alpha$  blockade has been observed in several animal models. Studies in diabetic animals with nephropathy showed that supplementation with vitamin E via reduced production of transforming growth factor-beta reduced urinary isoprostane level and improved proteinuria and blood urea nitrogen level [36]. In spontaneously animal models of hypertension and angiotensin II-induced hypertension, both with elevated plasma isoprostane level, Tempol administration decreases renal isoprostane excretion and lowers blood pressure [4,37].

These beneficial effects observed by isoprostane blockade (and increase in NO availability) can prevent sodium overload, hypertension, chronic fibrosis, and end-organ damage. Clinical evaluations are under review to estimate the beneficial effect of isoprostane blockade (NCT03358524, NCT01125501, and NCT00552227).

Although the results of this report can explain important pathophysiological conditions, it has some limitations. One important limitation is the H89, despite being the most commonly used PKA inhibitor, might have nonspecific inhibitory effects.

In conclusion, 8-iso-PGF2 $\alpha$  can increase J<sub>Cl</sub> at the kidney cTAL via stimulation of PKA even when J<sub>Cl</sub> is inhibited by NO. This increase of J<sub>Cl</sub> is associated with a higher cAMP tissue concentration, which cannot be inhibited by NO, indicating the importance of 8-iso-PGF2 $\alpha$  as a therapeutic intervention in kidney repair and functional restoration to regulate fluid and electrolyte balance.

### Conflicts of interest

All authors have no conflicts of interest to declare.

## Funding

This work was partially supported by Grant PIP 6525 from CONICET, Argentina, and by grant from the J. Robert Cade Foundation (No. L#003).

## Acknowledgments

Néstor H. García acknowledges the support from the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

## Authors' contributions

Conceptualization, Funding acquisition: LIJ, NHG  
Investigation, Investigation, Methodology: PDC, STB, GBS  
Data curation, Formal analysis: GBS, LIJ, NHG  
Writing—original draft: GBS, NHG  
Writing—review & editing: PDC, STB, LIJ, EIOA, NHG  
All authors read and approved the final manuscript.

## ORCID

Pablo D. Cabral, <https://orcid.org/0000-0001-5400-4291>  
Guillermo B. Silva, <https://orcid.org/0000-0001-9168-3369>  
Sandra T. Baigorria, <https://orcid.org/0000-0002-5447-0943>  
Luis I. Juncos, <https://orcid.org/0000-0002-3840-0419>  
Ebenezer I. O. Ajayi, <https://orcid.org/0000-0001-6348-5840>  
Néstor H. García, <https://orcid.org/0000-0002-9057-9030>

## References

1. Cabral PD, Silva GB, Baigorria ST, Juncos LA, Juncos LI, García NH. 8-iso-prostaglandin-F2 $\alpha$  stimulates chloride transport in thick ascending limbs: role of cAMP and protein kinase A. *Am J Physiol Renal Physiol* 2010;299:F1396–F1400.
2. Wang Y, Chen AF, Wang DH. Enhanced oxidative stress in kidneys of salt-sensitive hypertension: role of sensory nerves. *Am J Physiol Heart Circ Physiol* 2006;291:H3136–H3143.
3. Vara-Messler M, Mukdsi JH, Osieki NI, et al. Eicosapentaenoic acid prevents salt sensitivity in diabetic rats and decreases oxidative stress. *Nutrition* 2020;72:110644.
4. Reckelhoff JF, Zhang H, Srivastava K, Roberts LJ 2nd, Morrow JD, Romero JC. Subpressor doses of angiotensin II increase plasma F(2)-isoprostanes in rats. *Hypertension* 2000;35(1 Pt 2):476–479.
5. Ren Y, D'Ambrosio MA, Garvin JL, Carretero OA. Angiotensin II enhances connecting tubule glomerular feedback. *Hypertension* 2010;56:636–642.
6. Ortiz PA, Garvin JL. Autocrine effects of nitric oxide on HCO<sub>3</sub>(-) transport by rat thick ascending limb. *Kidney Int* 2000;58:2069–2074.
7. Gonzalez-Vicente A, Saez F, Monzon CM, Asirwatham J, Garvin JL. Thick ascending limb sodium transport in the pathogenesis of hypertension. *Physiol Rev* 2019;99:235–309.
8. García NH, Plato CF, Stoos BA, Garvin JL. Nitric oxide-induced inhibition of transport by thick ascending limbs from Dahl salt-sensitive rats. *Hypertension* 1999;34:508–513.
9. Garvin JL, Burg MB, Knepper MA. Ammonium replaces potassium in supporting sodium transport by the Na-K-ATPase of renal proximal straight tubules. *Am J Physiol* 1985;249(5 Pt 2):F785–F788.
10. García NH, Plato CF, Garvin JL. Fluorescent determination of chloride in nanoliter samples. *Kidney Int* 1999;55:321–325.
11. Plato CF, Stoos BA, Wang D, Garvin JL. Endogenous nitric oxide inhibits chloride transport in the thick ascending limb. *Am J Physiol* 1999;276:F159–F163.
12. Ortiz PA, Garvin JL. NO Inhibits NaCl absorption by rat thick ascending limb through activation of cGMP-stimulated phosphodiesterase. *Hypertension* 2001;37(2 Pt 2):467–471.
13. Morrow JD, Minton TA, Mukundan CR, et al. Free radical-induced generation of isoprostanes in vivo: evidence for the formation of D-ring and E-ring isoprostanes. *J Biol Chem* 1994;269:4317–4326.
14. Morla L, Edwards A, Crambert G. New insights into sodium transport regulation in the distal nephron: role of G-protein coupled receptors. *World J Biol Chem* 2016;7:44–63.
15. Praticó D, FitzGerald GA. Generation of 8-epiprostaglandin F<sub>2</sub> $\alpha$  by human monocytes: discriminate production by reactive oxygen species and prostaglandin endoperoxide synthase-2. *J Biol Chem* 1996;271:8919–8924.
16. Schuster VL. Molecular mechanisms of prostaglandin transport. *Annu Rev Physiol* 1998;60:221–242.
17. Badr KE, Abi-Antoun TE. Isoprostanes and the kidney. *Antioxid Redox Signal* 2005;7:236–243.
18. Salem MM. Pathophysiology of hypertension in renal failure. *Semin Nephrol* 2002;22:17–26.
19. Bauer J, Ripperger A, Frantz S, Ergün S, Schwedhelm E, Bendorff RA. Pathophysiology of isoprostanes in the cardiovascular system: implications of isoprostane-mediated thromboxane A<sub>2</sub> receptor activation. *Br J Pharmacol* 2014;171:3115–3131.

20. Caceres PS, Ares GR, Ortiz PA. cAMP stimulates apical exocytosis of the renal Na(+)-K(+)-2Cl(-) cotransporter NKCC2 in the thick ascending limb: role of protein kinase A. *J Biol Chem* 2009;284:24965–24971.
21. Rieg T, Kohan DE. Regulation of nephron water and electrolyte transport by adenylyl cyclases. *Am J Physiol Renal Physiol* 2014;306:F701–F709.
22. Carlström M, Brown RD, Sällström J, et al. SOD1 deficiency causes salt sensitivity and aggravates hypertension in hydronephrosis. *Am J Physiol Regul Integr Comp Physiol* 2009;297:R82–92.
23. Majid DS, Nishiyama A. Nitric oxide blockade enhances renal responses to superoxide dismutase inhibition in dogs. *Hypertension* 2002;39:293–297.
24. Kopkan L, Majid DS. Superoxide contributes to development of salt sensitivity and hypertension induced by nitric oxide deficiency. *Hypertension* 2005;46:1026–1031.
25. Ortiz PA, Garvin JL. Superoxide stimulates NaCl absorption by the thick ascending limb. *Am J Physiol Renal Physiol* 2002;283:F957–F962.
26. Ortiz MC, Sanabria E, Manriquez MC, Romero JC, Juncos LA. Role of endothelin and isoprostanes in slow pressor responses to angiotensin II. *Hypertension* 2001;37(2 Pt 2):505–510.
27. Ares GR, Caceres P, Alvarez-Leefmans FJ, Ortiz PA. cGMP decreases surface NKCC2 levels in the thick ascending limb: role of phosphodiesterase 2 (PDE2). *Am J Physiol Renal Physiol* 2008;295:F877–F887.
28. Ares GR. cGMP induces degradation of NKCC2 in the thick ascending limb via the ubiquitin-proteasomal system. *Am J Physiol Renal Physiol* 2019;316:F838–F846.
29. Vazzana N, Santilli F, Lattanzio S, et al. Determinants of thromboxane biosynthesis in patients with moderate to severe chronic kidney disease. *Eur J Intern Med* 2016;33:74–80.
30. Dimitrakov D, Kumchev E, Lyutakova E, Ledzhev I, Nikolov D, Tsekov V. Cyclic adenosine monophosphate (cAMP) levels in predialysis patients with chronic renal failure. *Folia Med (Plovdiv)* 1997;39:29–33.
31. Baylis C. Nitric oxide deficiency in chronic kidney disease. *Am J Physiol Renal Physiol* 2008;294:F1–F9.
32. Krishnan SM, Kraehling JR, Eitner F, Bénardeau A, Sandner P. The impact of the nitric oxide (NO)/soluble guanylyl cyclase (sGC) signaling cascade on kidney health and disease: a preclinical perspective. *Int J Mol Sci* 2018;19:1712.
33. Kaur J, Young BE, Fadel PJ. Sympathetic overactivity in chronic kidney disease: consequences and mechanisms. *Int J Mol Sci* 2017;18:1682.
34. Remuzzi G, Perico N, Macia M, Ruggenenti P. The role of renin-angiotensin-aldosterone system in the progression of chronic kidney disease. *Kidney Int Suppl* 2005;(99):S57–S65.
35. Larivière R, Lebel M. Endothelin-1 in chronic renal failure and hypertension. *Can J Physiol Pharmacol* 2003;81:607–621.
36. Montero A, Munger KA, Khan RZ, et al. F(2)-isoprostanes mediate high glucose-induced TGF-beta synthesis and glomerular proteinuria in experimental type I diabetes. *Kidney Int* 2000;58:1963–1972.
37. Schnackenberg CG, Wilcox CS. Two-week administration of tempol attenuates both hypertension and renal excretion of 8-Iso prostaglandin f2alpha. *Hypertension* 1999;33(1 Pt 2):424–428.