

Henry Ford Health System

Henry Ford Health System Scholarly Commons

Diagnostic Radiology Articles

Diagnostic Radiology

6-1-2014

Dynamic contrast enhanced MRI parameters and tumor cellularity in a rat model of cerebral glioma at 7 T

Madhava P. Aryal

Henry Ford Health System

Tavarekere N. Nagaraja

Henry Ford Health System, TNagara1@hfhs.org

Kelly A. Keenan

Henry Ford Health System, kkeenan1@hfhs.org

Hassan Bagher-Ebadian

Henry Ford Health System, HBAGHER1@hfhs.org

Swayamprava Panda

Henry Ford Health System, spanda1@hfhs.org

See next page for additional authors

Follow this and additional works at: https://scholarlycommons.henryford.com/radiology_articles

Recommended Citation

Aryal MP, Nagaraja TN, Keenan KA, Bagher-Ebadian H, Panda S, Brown SL, Cabral G, Fenstermacher JD, and Ewing JR. Dynamic contrast enhanced MRI parameters and tumor cellularity in a rat model of cerebral glioma at 7 T. *Magn Reson Med* 2014; 71(6):2206-2214.

This Article is brought to you for free and open access by the Diagnostic Radiology at Henry Ford Health System Scholarly Commons. It has been accepted for inclusion in Diagnostic Radiology Articles by an authorized administrator of Henry Ford Health System Scholarly Commons.

Authors

Madhava P. Aryal, Tavarekere N. Nagaraja, Kelly A. Keenan, Hassan Bagher-Ebadian, Swayamprava Panda, Stephen L. Brown, Glauber Cabral, Joseph D. Fenstermacher, and James R. Ewing

Dynamic Contrast Enhanced MRI Parameters and Tumor Cellularity in a Rat Model of Cerebral Glioma at 7 T

Madhava P. Aryal,^{1,2} Tavarekere N. Nagaraja,³ Kelly A. Keenan,³ Hassan Bagher-Ebadian,^{1,4} Swayamprava Panda,² Stephen L. Brown,⁵ Glauber Cabral,² Joseph D. Fenstermacher,^{3,6} and James R. Ewing^{1,2*}

Purpose: To test the hypothesis that a noninvasive dynamic contrast enhanced MRI (DCE-MRI) derived interstitial volume fraction (v_e) and/or distribution volume (V_D) were correlated with tumor cellularity in cerebral tumor.

Methods: T_1 -weighted DCE-MRI studies were performed in 18 athymic rats implanted with U251 xenografts. After DCE-MRI, sectioned brain tissues were stained with Hematoxylin and Eosin for cell counting. Using a Standard Model analysis and Logan graphical plot, DCE-MRI image sets during and after the injection of a gadolinium contrast agent were used to estimate the parameters plasma volume (v_p), forward transfer constant (K^{trans}), v_e , and V_D .

Results: Parameter values in regions where the standard model was selected as the best model were: (mean \pm S.D.): $v_p = (0.81 \pm 0.40)\%$, $K^{trans} = (2.09 \pm 0.65) \times 10^{-2} \text{ min}^{-1}$, $v_e = (6.65 \pm 1.86)\%$, and $V_D = (7.21 \pm 1.98)\%$. The Logan-estimated V_D was strongly correlated with the standard model's $v_p + v_e$ ($r = 0.91$, $P < 0.001$). The parameters, v_e and/or V_D , were significantly correlated with tumor cellularity ($r \geq -0.75$, $P < 0.001$ for both).

Conclusion: These data suggest that tumor cellularity can be estimated noninvasively by DCE-MRI, thus supporting its utility in assessing tumor pathophysiology. **Magn Reson Med** 71:2206–2214, 2014. © 2013 Wiley Periodicals, Inc.

Key words: dual-echo gradient-echo sequence; longitudinal relaxation rate; DCE-MRI; Logan plot; interstitial volume; distribution volume; tumor cellularity

Dynamic contrast enhanced MRI (DCE-MRI), because it is minimally invasive and free of radiation exposure, is becoming the technique of choice for assessing cerebral vascular permeability (1). A series of T_1 -weighted images are acquired before, during, and after the injection of a

contrast agent (CA), and a pharmacokinetic model is used to estimate blood-to-brain transfer parameters.

The microvascular leakage of a CA can be parameterized by its plasma volume (v_p), forward transvascular transfer constant (the rate of vessel-to-tissue leakage of the CA) K^{trans} , and, if the leakage occurs swiftly enough, by the CA's reverse transfer constant (the rate of tissue-to-vessel CA return), k_{ep} (2–6). If there are no active transport mechanisms, and no trapping of CA in either vascular or extravascular compartments, and when all three parameters (v_p , K^{trans} , and k_{ep}) can be stably estimated, the derivation of the interstitial volume fraction, v_e , then follows the relationship: $v_e = \frac{K^{trans}}{k_{ep}}$.

While the use of graphical methods for estimating model parameters has decreased in proportion to the availability of modern computers, graphical methods' utility for examining model plausibility remains. Two commonly applied graphical methods in concentration-time studies are the Patlak plot (2), and Logan plot (7,8). The Patlak plot was introduced in MRI (9) to estimate K^{trans} in a rat model of transient cerebral ischemia, and the extended Patlak method (3) has been used in a rat model of cerebral tumor (9). This latter method, or the equivalent extended Tofts model (4), also known as the standard model (SM), is now frequently used for the estimation of the vascular parameters v_p , K^{trans} , and v_e in DCE-MRI. However, DCE-MRI-derived v_e has not yet been evaluated as a true measure of interstitial space in brain tumor, nor has its relationship to tumor cellularity been examined. As for the Logan plot, we know of no instance of its use in DCE-MRI for the estimation of distribution volume (i.e., plasma volume, v_p , plus interstitial volume, v_e , expressed as a fraction), and no examination of its relationship to tumor cellularity. In this study, we introduce the Logan plot in DCE-MRI for the estimation of distribution volume (V_D) in an experimental cerebral tumor and assess its relationship to tumor cellularity.

Several studies have reported the correspondence of DCE-MRI-derived vascular parameters with histological findings, including tumor extracellular space, microvascular density, vascular endothelial growth factor, and tumor grade (10–12). In a study that is particularly relevant to the considerations of this study, Aref et al. demonstrated a strong correlation between a DCE-MRI-derived estimate of extracellular space volume fraction, and histologically determined extracellular space in a rat model of chemically induced mammary tumors (10).

Because tumor cellularity, i.e., the number of cells in a given volume of tumor tissue, is an important factor for proliferative activity (13), a comparison between noninvasive MRI measures and histology is a significant test

¹Department of Physics, Oakland University, Rochester, Michigan, USA.

²Department of Neurology, Henry Ford Hospital, Detroit, Michigan, USA.

³Department of Anesthesiology, Henry Ford Hospital, Detroit, Michigan, USA.

⁴Department of Radiology, Henry Ford Hospital, Detroit, Michigan, USA.

⁵Department of Radiation Oncology, Henry Ford Hospital, Detroit, Michigan, USA.

⁶Department Neurological Surgery, SUNY Stony Brook Medical University, Stony Brook, New York, USA.

Grant sponsor: National Institutes of Health; MRI Biomarkers of Response in Cerebral Tumors; Grant number: R01 CA135329-01.

*Correspondence to: James R. Ewing, Ph.D., Department of Physics, Oakland University, Rochester, MI. E-mail: jre@neurnis.neuro.hfh.edu

Received 11 December 2012; revised 8 May 2013; accepted 15 June 2013
DOI 10.1002/mrm.24873

Published online 22 July 2013 in Wiley Online Library (wileyonlinelibrary.com).

for validation of the clinical utility of DCE-MRI-derived parameters. Thus, the objective of this study was to test the hypothesis that tumor cellularity and a DCE-MRI-derived estimate of v_e and/or V_D were correlated in experimental cerebral tumor. In the course of examining the evidence, a number of other measures were examined, but the relation between tumor cellularity and the DCE-MRI estimated v_e and/or V_D was the main consideration of this analysis.

THEORY

The Patlak Graphical Analysis and Its Relationship to the SM

A simplified matrix-based model mathematically equivalent to the Crone-Renkin single-capillary model with no extra- to intra-vascular efflux was introduced by Patlak et al. (2). Subsequently, the original Patlak model was extended to allow for efflux from interstitial space to the vasculature (3). The extended Patlak model is identical to the extended Tofts, or consensus model (4), and is herein referred to as the SM. The SM describes the concentration of CA in tissue after intravenous administration:

$$C_t(t) = K^{\text{trans}} \int_0^t C_p(\tau) e^{-k_{\text{ep}}(t-\tau)} d\tau + v_p C_p(t), \quad [1]$$

where t is time, $C_p(t)$ and $C_t(t)$ are the time-varying plasma and tissue concentrations of CA, respectively, K^{trans} is the forward transfer constant of CA between blood plasma and interstitial space, k_{ep} is the reverse transfer rate constant from the interstitial space to the vascular compartment, and v_p is the vascular plasma volume fraction.

If it is assumed that the change in longitudinal relaxation rate ($R_1 = 1/T_1$) is proportional to tissue concentration of the CA (14), the observation equation equivalent to the SM equation is:

$$(1 - \text{Hct}) \Delta R_{1t}(t) = K^{\text{trans}} \int_0^t \Delta R_{1a}(\tau) e^{-k_{\text{ep}}(t-\tau)} d\tau + v_p \Delta R_{1a}(t), \quad [2]$$

where ΔR_{1a} and ΔR_{1t} refer to the subtraction of the pre-contrast relaxation rate from its post-contrast value at artery and tissue, respectively, and Hct is the hematocrit at the level of the microvasculature. In what follows, Hct was assumed to be 0.45 (15), which may uniformly bias the results presented below, but will not affect the correlations presented.

Plotting the ratio $\frac{(1-\text{Hct})\Delta R_{1t}(t)}{\Delta R_{1a}(t)}$ versus $\frac{\int_0^t \Delta R_{1a}(\tau) e^{-k_{\text{ep}}(t-\tau)} d\tau}{\Delta R_{1a}(t)}$, with k_{ep} optimized to provide the best fit (in the least-squares sense) of the model to the data yields the extended Patlak plot, with a slope of K^{trans} and an ordinate intercept of v_p . In the tumor, because, within the duration of the DCE experiment, the CA leaks from the microvasculature at a high enough rate to begin to equilibrate between the intra- and extra-vascular space, k_{ep} can also be estimated. Because this model fits three parameters: v_p , K^{trans} , and k_{ep} , it is called model 3. We

emphasize that model 3 results are identical to the results produced by the SM, but with the proviso that the SM has been tested against a reduced model, and thus more reliably represents the data. Note that, because an estimate of v_e requires the ratio $K^{\text{trans}}/k_{\text{ep}}$, DCE-MRI estimates of interstitial volume can be reliably performed only in model 3 regions. The same goes for Logan analysis estimates of V_D (see below).

In some regions with CA leakage, but no measurable back-flux to the vasculature (i.e., $k_{\text{ep}} = 0$), only two parameters, K^{trans} and v_p , can be estimated. The model in this case has two parameters and is named model 2. In normal brain, there is usually little or no measurable leakage of CA from the microvasculature. In that case, $K^{\text{trans}} = k_{\text{ep}} = 0$; the one parameter to be estimated is v_p and is accordingly identified as model 1. Model selection proceeded as previously described (6): for each voxel, the preferred model was selected by the application of an F -test criterion.

The Logan Plot

The tissue and plasma concentration of the CA as a function of time is (7,8);

$$\frac{\int_0^t C_t(\tau) d\tau}{C_t(t)} = V_D \frac{\int_0^t C_p(\tau) d\tau}{C_t(t)} + \text{Const} \quad [3]$$

where $C_p(t)$ and $C_t(t)$ are the plasma and tissue concentrations of CA, respectively, V_D is the distribution volume, and, if an equilibrium has been established between the vascular compartment and the tissue compartment, Const is a constant. We note that the Logan plot fits a different portion of the data in DCE-MRI studies than the SM: the SM depends on changes in the response function (relative to the input function) to estimate the rate parameters K^{trans} and k_{ep} , while the Logan analysis demands that the response function does not change relative to the input function. Thus, the two analyses fit much different portions of the input-response data, albeit they both require a measure of the data from time $t = 0$. Note that, even if k_{ep} is not explicitly estimated (Eq. [2]), but expressed as $\frac{K^{\text{trans}}}{v_e}$ in the exponent of the kernel, it is only through dynamic changes that the term can be estimated.

Because the change in longitudinal relaxation rate is assumed to be approximately proportional to tissue concentration of the CA, the observation equation for the Logan plot becomes:

$$(1 - \text{Hct}) \frac{\int_0^t \Delta R_{1t}(\tau) d\tau}{\Delta R_{1t}(t)} = V_D \frac{\int_0^t \Delta R_{1a}(\tau) d\tau}{\Delta R_{1a}(t)} + \text{Const} \quad [4]$$

where $\Delta R_{1a}(t)$ and $\Delta R_{1t}(t)$ refers to the subtraction of the pre-contrast relaxation rate from its post-contrast value at artery and tumor region, respectively, as a function of time, and Hct is the hematocrit.

If there is a time after which the plot of $(1 - \text{Hct}) \frac{\int_0^t \Delta R_{1t}(\tau) d\tau}{\Delta R_{1t}(t)}$ versus $\frac{\int_0^t \Delta R_{1a}(\tau) d\tau}{\Delta R_{1a}(t)}$ yields a straight line, the slope of the fitted straight line gives the distribution

volume. Once again, the graphical nature of this statement is central to the choice of model. Generally in this article, data were examined visually for conformance to the model before selection of the point for starting the fit of the Logan plot. This procedure can be automated if it is to be performed on a voxel-by-voxel basis, but a preliminary visual examination of grouped data should be performed before moving on to a voxel-by-voxel procedure.

ΔR_1 Calculation Theory

Under the conditions of the experiment, i.e., using a dual-echo gradient echo sequence (the mgems sequence in the Agilent VNMRj library), and assuming that the flip angle is a known constant across the brain of the rodent, an estimate of the time trace of ΔR_1 can be calculated from the following:

$$S(t) = \frac{\varepsilon M_0 \sin(\theta)(1 - e^{-TRR_1(t)})e^{-TER_2^*(t)}}{1 - \cos(\theta)e^{-TRR_1(t)}}, \quad [5]$$

where $S(t)$ the signal intensity at time t , ε is the detection efficiency, M_0 is the equilibrium magnetization of the protons, θ is the tip-angle, TR is the repetition time between pulses, TE is the echo time (the time between the center of the excitation pulse and the readout gradient), and $R_2^*(t)$ is the transverse relaxation rate. Equation [[5]] can be solved for $R_1(t)$:

$$R_1(t) = \frac{1}{TR} \ln \left\{ \frac{1 - \left(\frac{S(t)\cos(\theta)}{\varepsilon M_0 e^{-TER_2^*} \sin(\theta)} \right)}{1 - \left(\frac{S(t)}{\varepsilon M_0 e^{-TER_2^*} \sin(\theta)} \right)} \right\} \quad [6]$$

METHODS

These studies were conducted with the approval of the Henry Ford Hospital Institutional Animal Care and Use Committee.

Animal Preparation

U251 Tumor Induction

Athymic rats were inoculated intracerebrally as follows: animals were anesthetized with 80 mg/kg ketamine and 15 mg/kg xylazine i.m. The surgical zone was swabbed with Betadine solution, the eyes coated with Lacri-lube, and the head immobilized in a small animal stereotactic device (Kopf, Cayunga, CA). After draping, a 1 cm incision was made 2 mm right of the midline and the skull was exposed. A burr hole was drilled 3.5 mm to the right of bregma, taking care not to penetrate the dura. A #2701 10 μ L Hamilton syringe with a 26 s gauge needle containing U251MG tumor cells freshly harvested from log phase growth (5×10^5 in 10 μ L of phosphate buffered solution (PBS)) was lowered to a depth of 3.0 mm and then raised back to a depth of 2.5 mm. Cells were then injected at a rate of 0.5 μ L/10 s until the entire volume was injected. Tumors in animals implanted following this technique grow to about 5 mm diameter by 21 days post-implantation. On or a few days after day 18, each

animal was anesthetized (isoflurane 4% for knockdown, 0.75–1.5% for maintenance, balance $N_2O:O_2 = 2:1$, and allowed to spontaneously respire). A tail vein was cannulated for the administration of CA. Body temperature was maintained constant (37°C) with a warm air supply monitored via an intrarectal type T thermocouple.

MRI Study

DCE-MRI studies were performed using a 7 T magnet in 18 athymic rats at 18 ± 3 days after tumor implantation. The image sets were acquired on a DirectDrive Varian (now Agilent, Santa Clara, CA) 21 cm bore system with a 32×32 mm field of view. Coils were a Bruker-supplied volume resonator and a Bruker 2 cm surface coil for rat brain imaging. High-resolution T_1 -weighted spin-echo images were acquired pre- and post-CA (flip angle (FA) = 90°, 180°, matrix 256×192 , 27 slices, 0.5 mm thickness, number of echoes (NE) = 1, number of averages (NA) = 4, TE/TR = 16/800 ms). Before the DCE-MRI sequence, and immediately after, two Look-Locker sequences (LL) sequences were run so that a voxel-by-voxel estimate of T_1 in the tissue could be made pre- and post-CA administration. LL sequence parameters were as follows: FA = 15°, matrix 128×64 , five 2.0 mm slices, no gap. NE = 24 inversion-recovery echoes on 50 ms intervals, TE/TR = 4.0/2000 ms.

The DCE-MRI sequence was a dual-echo gradient-echo sequence, the “mgems” sequence in the Agilent Vnmrj library. It acquired 150 image sets at 4.0 s intervals: (FA = 27°, matrix = 128×64 , three 2.0 mm slices, NE = 2, NA = 1, TE1/TE2/TR = 2.0/4.0/60 ms). At image 15 of the dual-echo gradient-echo sequence Bolus injection of the CA (Magnevist, Bayer Healthcare Pharmaceuticals, Wayne, NJ), 0.25 mmol/kg at undiluted concentration, no flush, was performed by hand push, followed by a slight draw-back. Following the post-contrast LL sequence, a Pulsed Gradient Spin-Echo DWI sequence (matrix 128×64 , thirteen 1 mm slices, TR = 1500 ms, TE = 40 ms, NE = 1, b -values = 0, 1217 s/mm^2 , gradient amplitude = 1.07 mT/cm, gradient duration = 10 ms) was run to generate a parametric map of apparent diffusion coefficient (ADC). These studies carried over from a previous study that used two CAs of different weights (Magnevist, then Gd-labeled albumin), with the ADC studies conducted after the first study in order to save time and allow for partial clearance of the first CA. Thus, there may be some small bias in the results (16). However, because all the studies were conducted in the same manner, and the results examined through correlation, we judged that inferences drawn from this study might still be useful.

Histology

At sacrifice, animals were deeply anesthetized with ketamine/xylazine and transcardiacally perfused with saline followed by 4% paraformaldehyde. After carefully removing the brains from the skull, they were post-fixed in same fixative overnight. Coronal sections grossly containing and adjacent to the tumor were obtained using a rat-brain matrix (Activational Systems, Inc., Warren, MI). The brain tissue was processed using a VIP Tissue Tek

Processing center, before embedding. Seven micrometer sections were cut and placed on Superfrost Plus (Fisher Scientific) slides and stained with Hematoxylin and Eosin (H&E) for evaluation of tumor regions of interest (ROIs). Additional sections were obtained for immunohistochemistry staining. Images were collected using a Nikon Eclipse E800 Microscope equipped with ACT1C software.

H&E stained slides were used for cell density measurements. ROIs were chosen in the tumor interior and in the tumor periphery. The area of the field of view in each ROI was 4000 μm by 3000 μm . A comparative ROI was chosen within the contralateral hemisphere, usually within the caudate putamen. Both 1 \times low magnification of the entire coronal section and 10 \times higher magnification images were collected. Images were imported into ImageJ (NIH, version 1.43u), and converted to 8 bits for analysis. A thresholding range of 133–193 was used, depending on the contrast of the H&E staining. The numbers of nuclei were manually counted in a sampling of regions and compared with the automated count to ensure accuracy. The particle analysis function used was set to collect particles in the range of 15–70 μm^2 . This was used to count cell nuclei and, therefore, used as the basis for the cell numbers measures.

Arterial Input Function

Because of the difficulty of directly estimating the arterial input function from MRI data, a radiotracer assay-based input function obtained from a previous investigation (17) was used in the dynamic MRI studies. Nagaraja et al. (17) have shown that, starting at about two minutes after injection of CA, the radiological and MRI measures of blood concentrations of CA track each other very well. In a number of studies, we examined the tissue response in a brain area where minimal leakage was known to occur. It was clear from an examination of the time trace of ΔR_1 that the early portion of the tissue response curve did not resemble that of an input function, presumably because of the dispersion in time of the CA arriving at, and passing through, the vascular system of the tissue, and also possibly because of limited water exchange across the blood-brain barrier. However, during the latter two-thirds of the study, the tissue response and the radiological standard curve we used appeared to agree, in line with Nagaraja's observations. Accordingly, the latter two-thirds of the time trace of the radiological curve and the tissue trace of $\Delta R_1(t)$ in the caudate putamen were integrated and used to scale the radiological curve. It was assumed that: (1) no vascular leakage of CA occurred in the caudate putamen region contralateral to the tumor, and (2) the plasma volume of the caudate putamen was 1% (18). The integrated area of the radiotracer arterial input function was scaled so that its area was 100 times that of the integrated area of the average value of $\Delta R_1(t)$ in the caudate putamen of the opposite hemisphere. The 1% figure adopted was larger than Bereczki's estimates (0.8–0.9%, Table 5 of the Bereczki paper (18)) because that work, as the authors noted, estimated only plasma volumes in vessels below 50 μm . All the volumes estimated in this article are relative to the

estimate of v_p in the caudate putamen; we took the value of 1% so that relative volumes measured elsewhere could be easily understood in relation to caudate putamen plasma volumes. This method is a combination of the reference region approach of Yankeelov et al. (19) and the original Tofts approach that used a group-averaged input function (20).

Parameter Estimation—Computational and Statistical Methods

The analysis of the LL data followed that of Gelman et al. (21). Briefly, because it was determined that, for a typical study, the flip angle, θ , did not vary by more than a few percent across a rat brain, a whole-brain estimate of θ was first performed by using a boxcar-smoothed set of inversion-recovery images and using a Simplex least-squares minimum finder in three parameters to freely fit ϵM_0 , T_1 , and θ voxel-by-voxel in the smoothed images (this ϵ was not the same as that of the DCE-MRI sequence). The flip angle was then fixed at the mean of its global estimate for the brain and a Simplex minimum finder in two parameters was used in the unsmoothed data to produce a voxel-by-voxel estimate of T_1 and ϵM_0 .

We note that the 2000 ms repetition time of the LL sequence was too short to allow the inverted magnetization to fully relax to its equilibrium value. An iterative procedure to estimate the equilibrium value of the magnetization (M_{ss} in Eq. [1] of Gelman et al. (21)) in the computational estimate of T_1 was used to adjust for the short TR's, allowing an unbiased estimator of T_1 to be produced.

Consider Eqs. [5] and [6]. Besides $R_1(t)$ and $R_2^*(t)$ which vary across the experiment, the parameters ϵM_0 and θ , which do not vary across the experiment, must be estimated. Our practice was to estimate R_2^* using the log ratio of the first and second echo, to fix the value of θ at 27°, and to use maps of R_1 constructed from LL maps, in combination with the amplitude of the DCE data at the beginning and end of the study, to estimate ϵM_0 , as described in the following paragraph.

The three unknowns in the DCE-MRI sequence (Eq. [5]) were $R_1(t)$, θ , and ϵM_0 . The measured quantity was $S(t)$. It was assumed that the tip angle θ was known. In the period before the administration of CA, using the voxel-by-voxel estimate of $R_1(t < t_{inj})$, produced by the prior LL measurements, where t_{inj} was the time of CA injection, the quantity ϵM_0 was estimated using Eq. [5]. Using the post-study estimates of R_1 and the last few points of the DCE-MRI data where the data was changing slowly, the quantity ϵM_0 was similarly estimated, and the average of the two estimates was taken. Assuming that the efficiency for detection did not change after the injection of CA, this allowed $R_1(t)$ to be estimated across time by means of Eq. [6].

The practice of calculating R_1 via LL sequences before and after the DCE-MRI study allowed ϵM_0 to be estimated under conditions of no CA, and approximately constant tissue CA concentration. This considerably stabilized the estimate of ϵM_0 , and thus of $R_1(t)$ because it fixed the beginning and end points of the $R_1(t)$ curve. The quantity $\Delta R_1(t)$ in Eq. [4] was generated by




FIG. 1. Left: High-resolution post-contrast T_1 -weighted image. Middle: Corresponding centrally located ADC maps. Right: H&E staining of a centrally located tissue slice approximating the position of the MRI image.

subtracting the pre-contrast estimate of $R_1(t)$ from the subsequent estimates of $R_1(t)$.

All results are reported as the mean \pm standard deviation. The standard error of the means in this sample of 18 studies is the standard deviation divided by 4.24.

The adjusted input function and tissue time trace of ΔR_1 were used to estimate pharmacokinetic parameters using the SM, as in Bagher-Ebadian et al. (6). For each voxel, data points were fitted to model 1, 2, and 3, an F -statistic was generated to select the preferred model, and the parameters available under that model were mapped. However, for the Logan plot, the adjusted input function and time trace of ΔR_1 in tumor tissue was used to generate the Logan graphical plot. The trend of the plot was then visually inspected for each animal individually to choose the starting time point for the straight line fitting of the plot, after which the slope of the fitted straight line was measured as an estimate of the distribution volume.

To estimate the DCE parameters for examining their correlation to tumor cellularity, a 5×4 pixel ROI was visually selected on parametric maps to correspond to the areas from the cell counting on histology. Because only one slice on MRI data set was used to make the comparison, anatomical markers such as lateral ventricles, coronal shape, and tumor area comparison (on high-resolution T_1 weighted image) were used to select the MRI maps and further pinpoint the histological (Histo) ROI onto the matching parametric maps. Hereafter, this 5×4 pixel ROI is named as the Histo sub-region.

Other Statistical Methods

The correlation of the Logan estimate of V_D , compared to the SM V_D , was examined by linear regression; their absolute estimates were assessed for differences via a paired t -test. Other relationships reported herein (e.g., the association between ADC and cell count) were assessed similarly.

RESULTS

Eighteen DCE-MRI animal studies and corresponding histological findings were available. All studies were the second study of a test-retest paradigm, with 24 h between studies. Test-retest findings are reported elsewhere (22). Mean tumor age was 18 ± 3 days. All tumors were seen clearly on post-contrast T_1 -weighted images, ADC maps, and H&E-stained tissue slides.

To introduce the findings, the data from an animal (#18) studied 21 days post-implantation will be presented. For the example animal, the mean ADC value (averaged from x , y , and z direction of the center image) was $(9.3 \pm 2.5) \times 10^{-4}$ mm²/s. A post-contrast T_1 -weighted image, the corresponding ADC map (averaged from x , y , and z direction) from the center slice, and an H&E-stained tissue slide approximating the position of the MR image of the example animal are shown in Figure 1.

Under the conditions of the experiment, the first pass of the CA produced a signal decrease due to T_2^* dephasing in many tissues, particularly in highly vascular regions of the tumor and in blood. Using the T_2^* -





FIG. 2. Parametric maps left to right: v_p , K^{trans} , v_e , and model selection. For the model selection map, white is model 3 acceptance, dark blue is model 2 acceptance, and light blue is model 1 acceptance.

FIG. 3. Left: H&E-stained tissue slide corresponding to parametric map. Right: Parametric map of v_e (only in the model 3 region can v_e be calculated). The white box drawn on the detail images shows the ROIs selected for cell count and v_e values.



corrected time trace of dual-echo gradient-echo data, the pre-contrast R_1 values, flip angle and repetition time, the time trace of ΔR_1 was calculated pixel-by-pixel across the entire image set. The time trace of ΔR_1 in turn was used to produce the parametric maps. The parametric maps for v_p , K^{trans} , v_e , and an F -test based model selection are shown in Figure 2. From the model selection map, we see that model 1 predominated in normal non-leaky brain tissue, showing little or no leakage of CA. Model 3 predominated in regions associated with a high density of tumor cells, as determined by examination of postmortem tissue slides. Model 2 bordered model 3 regions, showing leakage at reduced rates.

As noted earlier, v_e and V_D , estimates of interstitial space can only be measured when CA leaks from the capillary into the interstitial space in an amount sufficient to measure its back flux to the vasculature. Because the primary goal for this study was to study the relationship between tumor cellularity and DCE-MRI-derived estimates of interstitial space, only the data from model 3 regions were analyzed. The parameter values in model 3 regions varied considerably from animal to animal, with the mean values $v_p = (0.81 \pm 0.40)\%$, $K^{\text{trans}} = (2.09 \pm 0.65) \times 10^{-2} \text{ min}^{-1}$, and $v_e = (6.65 \pm 1.86)\%$.

Figure 3 shows the H&E-stained tissue slide with corresponding parametric map of v_e . The white box on the magnified image shows the Histo sub-region chosen for cell counts and parameter values, respectively. The mean parameter values estimated in the Histo sub-region were $v_p = (0.81 \pm 0.44)\%$, $K^{\text{trans}} = (2.08 \pm 1.45) \times 10^{-2} \text{ min}^{-1}$, and $v_e = (6.99 \pm 1.45)\%$.

For each time point, the average ΔR_1 across the points of the model 3 region and Histo sub-region was determined and used as a measure of the CA concentration for the purpose of producing a Logan plot. Figure 4(top) plots the data as an extended Patlak plot in the model 3 region; the slope of the fitted straight line gives K^{trans} . In this example, although the overall fit is excellent ($R^2 = 0.992$), there is a clear deviation from linearity in the early time points. The sources of this deviation are briefly considered in the Discussion.

Figure 4(bottom) plots the data as a Logan plot in the same region, with fitting for a straight line between abscissa values of 90:150 min. In that region, the graph was linear by inspection. The slope of the fitted line was 0.0897, equivalent to a distribution volume, V_D , of 8.97%.

For the sample of 18 animals in the study, the mean V_D measured in model 3 region was $(7.21 \pm 1.98)\%$, roughly half that of normal brain parenchyma (23). V_D estimated by two independent methods, SM and Logan analysis, was highly correlated ($r = 0.91$, $P < 0.001$), where, for the SM method, V_D was calculated as the sum of v_e and v_p . The mean SM V_D (7.46%) did not differ significantly from that of the Logan V_D (7.21%) ($t = 0.38$, $N = 18$, $P = 0.70$). This was a confirmatory result, because the two methods used primarily early data in the SM pharmacokinetic approach, and primarily late data in the Logan pharmacokinetic approach.

Cell counts from the Histo sub-region adjacent to the tumor core region varied from 751 to 3776 with mean (\pm standard deviation) 2442 ± 840 . Figure 5(left) shows a scatter plot of the SM estimated v_e versus tumor cellularity. Figure 5(right) shows the scatter plot of the Logan plot estimated V_D versus tumor cellularity. A linear regression of tumor cellularity with both v_e estimated via SM and V_D via Logan plot showed an inverse relationship, with a strong correlation ($r = -0.75$ for v_e and $r = -0.76$ for V_D , P -value < 0.001 for both). The mean SM estimated distribution volume (plasma volume plus interstitial volume) was also highly correlated with tumor cellularity ($r = -0.76$, P -value < 0.001). Thus, a measure of interstitial volume fraction (v_e) estimated from the SM, and distribution volume from both SM and Logan analyses were highly (negatively) correlated with the tumor cellularity.

The relationship of tumor cellularity and the ADC was also examined. A significant correlation with ADC ($r = 0.54$, P -value < 0.05) was observed. Notably, this relationship was not as strong as those of the DCE-MRI measures, although, because the ADC estimate occurred postcontrast, the result might be weakened.

DISCUSSION

In this study, we measured DCE-MRI parameters using a dual-echo gradient-echo sequence in a rat model of cerebral glioma at 7 T in order to establish their relationship to tumor cellularity. The mean DCE parameter values measured in model 3 regions and/or the Histo sub-regions were in agreement to those of similar measures in human glioblastoma using DCE-MRI (6), or CT perfusion (24), and in a rat model of 9L glioma using DCE-MRI (25). In this latter reference, however, estimates of




FIG. 4. Top: Patlak graphical analysis of the data in the Histo sub-region. The solid line shows the linear fit of the data points; the slope of the line yields the forward transfer constant, K^{trans} . Bottom: Logan graphical analysis of the data in the same sub-region. The solid line shows the linear fit of the data in the range 90:150, the slope of the line yields an estimate of distribution volume of about 9%.

v_e were much higher than our estimates. Whether the 9L glioma differs so radically from the U251 model used herein, or the differing analyses employed produce much different estimates of k_{ep} , remains to be determined. One possibility is that, because all the volume estimates were indexed to the plasma volume of the caudate putamen, the 1% value we used for plasma volume fraction in that tissue was too small.

The mean values in the entire model 3 region and the Histo sub-region did not differ for v_p and K^{trans} . However for interstitial volume, the mean values in the Histo sub-region were slightly elevated compared with model 3 region; the difference in mean values between two regions was not significantly different from zero ($P = 0.57$). As noted, the Histo sub-region was typically selected nearer the tumor core region; that might be a reason for the higher estimates of v_e in the Histo sub-region compared with the model 3 region as a whole. In

a study in a rat model of 9L glioma, v_e was also found to be elevated in the tumor core region (25). This result supports our higher estimates of mean v_e and/or V_D values in the Histo sub-region comparison with the model 3 region.

Although a number of DCE-MRI studies in glioma have evaluated blood volume and permeability, only a few have estimated interstitial volume and its relation to histological findings. In a study in human glioma, estimation of interstitial volume was helpful in differentiating gliomas from other brain tumors; increasing interstitial volume tended to correlate with increasing tumor grades (12). In a study of a rat model of chemically induced mammary tumors, Aref et al. demonstrated that interstitial volume fraction measured by DCE-MRI was correlated with histologically measured extracellular space fraction (10).

In another study in a rat model of 9L tumor, Chenevert et al. reported an increased volume of water in the interstitial compartment with decrease in cellularity following the chemotherapeutic (Carmustine) intervention (26). A strong correlation of mean ADC values with tumor cellularity ($r = 0.78$) was demonstrated, and also that those values were highly sensitive to changes in cellularity.




FIG. 5. Top: Scatter plot of Patlak plot estimates of interstitial volume versus tumor cellularity in the Histo sub-region. The solid line shows the linear fitting of the plot ($r = 0.75$, $P < 0.001$). Bottom: Scatter plot of Logan plot estimates of distribution volume (V_D) versus tumor cellularity measured in the Histo sub-region. Solid line shows the linear fitting of the plot ($r = 0.76$, $P < 0.001$).

This increase in interstitial volume could be due to vasogenic edema or shifts of intracellular water to the extracellular space associated with tumor cell necrosis. This result demonstrates the relationship of tumor cellularity with interstitial volume and ADC values; a decrease in cellularity causes the increase in the interstitial volume and ADC values.

This study has demonstrated the first use of a Logan plot in DCE-MRI, showing a strong correlation between tumor cell count and DCE-MRI derived interstitial volume and/or distribution volume estimated from two independent methods, SM and Logan analyses. The findings imply that the DCE-MRI estimated interstitial volume fraction and/or distribution volume can be used as a biomarker for tumor cellularity. They support not only the utility of DCE-MRI in assessing vascular permeability but also suggest that a measure of tumor cellularity is available for assessing and potentially classifying gliomas. Thus, DCE-MRI with appropriate model selection may be a useful tool for both the diagnosis and evaluation of therapeutic response in patients with gliomas.

Given that an estimate of V_D is available from the SM's estimate of $v_p + v_e$, one might reasonably ask "why generate a Logan plot of V_D ?" There are two arguments for the use of the Logan analysis. The first is that, in the event that a binding potential for a Gd-tagged agent that does bind to elements of the parenchyma (e.g., gadofosveset (27)) is needed, the operating characteristics of a compound that does not bind will be useful.

A second, and perhaps more relevant, support for Logan analysis when a nonbinding CA is an indicator is that the Logan estimate of V_D is likely to be more stable and less dependent on technique than that of the SM. In the SM, v_e is estimated from the ratio $K^{\text{trans}}/k_{\text{ep}}$, and is thus a derived parameter. Both of the parameters from which it is derived are rates: these cannot be estimated from equilibrium data, but must use those portions of the data that are dynamically changing with respect to the input function. Thus, when tissue in a DCE-MRI study is viewed as an input-response system, the response measures in the SM and the Logan graphical analysis fit much different portions of the response curves for their estimates of distribution volume, albeit they both require an integral of the input and response from time zero. In this presentation, it is gratifying that the two estimates are in good agreement, but we note the likelihood that SM estimates of distribution volume will be more dependent on high temporal resolution estimates of the input and tissue response function than will the Logan analysis. In support of this hypothesis, we note that, if the data of the example animal, with intrinsic 4 s time resolution, is binned into 12 s bins, the Logan estimate of V_D changes from 8.9% to 9.2%, while the SM estimate changes from 11% to 17%. However, because the operating characteristics of the Logan plot in DCE-MRI have not been explored, some caution should be exercised in assuming that it is less sensitive than the SM to temporal sampling rates.

We note also the evident deviation of the data from the straight-line extended Patlak fit in the early time points of Figure 4a (the extended Patlak approach is equivalent, as we have noted, to the SM). This is despite

an excellent overall fit of the model to the data ($R^2 = 0.992$). Excellent fits of this nature are standard in this approach; in the 18 studies, the lowest R^2 in model 3 grouped data was 0.972, and the group mean was 0.993. Nevertheless, for the summed data the fit is typically not perfect. As Figure 4a implies, there may be some additional systematic behavior in the first few minutes of tissue response that is not accounted for in the model.

The candidates for this systematic behavior are numerous. On the luminal side of the microvasculature, it may be that, in this particular animal, the radiological input function is not correct; it may be that the heterogeneity of the tumor in the model 3 region changes the arrival time and shape of the input function across the regions of the tumor, and thus, while the radiological input function is approximately correct, the input function at the various tumor regions is not; in a related hypothesis, it may be that the dispersion of the arterial input function due to flow changes the input at the tissue level; it may be that restricted water exchange in the microvasculature of the tumor changes the apparent longitudinal relaxivity of the tissue in a concentration-dependent manner (28–30). In the tissue, it may be that transport of the CA takes place at a relatively high rate, thus draining the tissue of CA and violating the model, which requires that CA, once it extravasates, is reabsorbed in the vasculature of the same voxel. All of these possibilities are plausible, and in fact may all contribute to the behavior of the signal in varying amounts.

The two effects that most probably account for the early deviation of the data from the linearity of the extended Patlak plot are transvascular water exchange effects and/or dispersion of the arterial input function due to flow. Unfortunately, it may not be possible to distinguish between the two.

As to whether water exchange affects the Logan analysis results, Li et al. (31) undertook a DCE-MRI study at 11.7 T in a rat model of cerebral glioma, with sequence parameters similar to this study, and included water exchange in their modeling. The dose of CA used in their study (0.5 mmol/kg) was twice that of ours, but they concluded that, while transvascular water exchange was observable, producing a negative bias in the estimate of plasma volume, transcytolemmal water exchange was not observable. That is to say, extravasated CA produced an approximately linear response in tissue ΔR_1 . Recent modeling that incorporates a full three-compartment water exchange model (32) supports this conclusion; for the relatively low values of K^{trans} observed in cerebral glioma, transcytolemmal water exchange has little effect. Thus, in the portion of the curve to which the Logan plot applies, when the arterial concentration has decreased, and equilibrated with the extravascular concentration of CA, water exchange effects are probably minimal. This tends to support the use of the Logan analysis in MRI, because it can eliminate one potential source of bias.

ACKNOWLEDGMENTS

The authors thank the following individuals: Jun Xu for assistance with animal preparations and histopathology,

Dr. Guangliang Ding for help in preparing MR sequences, and Dr. Robert Knight for help in running the MR sequences.

REFERENCES

- Hylton N. Dynamic contrast-enhanced magnetic resonance imaging as an imaging biomarker. *J Clin Oncol* 2006;24:3293-3298.
- Patlak CS, Blasberg RG, Fenstermacher JD. Graphical evaluation of blood-to-brain transfer constants from multiple-time uptake data. *J J Cereb Blood Flow Metab* 1983;3:1-7.
- Patlak C, Blasberg R. Graphical evaluation of blood to brain transfer constants from multiple time up take data. Generalizations. *J J Cereb Blood Flow Metab* 1985;5:584-590.
- Tofts PS, Brix G, Buckley DL, et al. Estimating kinetic parameters from dynamic contrast-enhanced T(1)-weighted MRI of a diffusable tracer: standardized quantities and symbols. *J Magn Reson Imaging* 1999;10:223-232.
- Ewing JR, Brown SL, Lu M, et al. Model selection in magnetic resonance imaging measurements of vascular permeability: Gadomer in a 9L model of rat cerebral tumor. *J Cereb Blood Flow Metab* 2006;26:310-320.
- Bagher-Ebadian H, Jain R, Nejad-Davarani SP, et al. Model selection for DCE-T1 studies in glioblastoma. *Magn Reson Med* 2012;68:241-251.
- Logan J, Fowler JS, Volkow ND, et al. Graphical analysis of reversible radioligand binding from time-activity measurements applied to [N-11C-methyl]-(-)-cocaine PET studies in human subjects. *J Cereb Blood Flow Metab* 1990;10:740-747.
- Logan J. Graphical analysis of PET data applied to reversible and irreversible tracers. *Nucl Med Biol* 2000;27:661-670.
- Ewing JR, Knight RA, Nagaraja TN, Yee JS, Nagesh V, Whitton PA, Li L, Fenstermacher JD. Patlak plots of Gd-DTPA MRI data yield blood-brain transfer constants concordant with those of ¹⁴C-sucrose in areas of blood-brain opening. *Magn Reson Med* 2003;50:283-292.
- Aref M, Chaudhari AR, Bailey KL, Aref S, Wiener EC. Comparison of tumor histology to dynamic contrast enhanced magnetic resonance imaging-based physiological estimates. *Magn Reson Imaging* 2008;26:1279-1293.
- Su MY, Cheung YC, Fruehauf JP, et al. Correlation of dynamic contrast enhancement MRI parameters with microvessel density and VEGF for assessment of angiogenesis in breast cancer. *J Magn Reson Imaging* 2003;18:467-477.
- Ludemann L, Grieger W, Wurm R, Budzisch M, Hamm B, Zimmer C. Comparison of dynamic contrast-enhanced MRI with WHO tumor grading for gliomas. *Eur Radiol* 2001;11:1231-1241.
- Kiss R, Dewitte O, Decaestecker C, Camby I, Gordower L, Delbecque K, Pasteels JL, Brotchi J, Salmon I. The combined determination of proliferative activity and cell density in the prognosis of adult patients with supratentorial high-grade astrocytic tumors. *Am J Clin Pathol* 1997;107:321-331.
- Cao Y, Shen Z, Chenevert TL, Ewing JR. Estimate of vascular permeability and cerebral blood volume using Gd-DTPA contrast enhancement and dynamic T₂*-weighted MRI. *J Magn Reson Imaging* 2006;24:288-296.
- Kuipers H, Brouwer T, Dubravcic-Simunjak S, Moran J, Mitchel D, Shobe J, Sakai H, Stray-Gundersen J, Vanhoutvin S. Hemoglobin and hematocrit values after saline infusion and tourniquet. *Int J Sports Med* 2005;26:405-408.
- Fitzek C, Mentzel HJ, Fitzek S, Sauner D, Kaiser WA, Reichenbach JR. Echoplanar diffusion-weighted MRI with intravenous gadolinium-DTPA. *Neuroradiology* 2003;45:592-597.
- Nagaraja TN, Karki K, Ewing JR, Divine GW, Fenstermacher JD, Patlak CS, Knight RA. The MRI-measured arterial input function resulting from a bolus injection of Gd-DTPA in a rat model of stroke slightly underestimates that of Gd-[(14)C]DTPA and marginally overestimates the blood-to-brain influx rate constant determined by Patlak plots. *Magn Reson Med* 2010;63:1502-1509.
- Bereczki D, Wei L, Otsuka T, Hans F-J, Acuff V, Patlak C, Fenstermacher J. Hypercapnia slightly raises blood volume and sizably elevates flow velocity in brain microvessels. *Am J Physiol* 1993;264:H1360-H1369.
- Yankeelov TE, Luci JJ, Lepage M, Li R, Debusk L, Lin PC, Price RR, Gore JC. Quantitative pharmacokinetic analysis of DCE-MRI data without an arterial input function: a reference region model. *Magn Reson Imaging* 2005;23:519-529.
- Tofts P, Kermode A. Measurement of the blood-brain barrier permeability and leakage space using dynamic MR Imaging. 1. Fundamental concepts. *Magn Reson Med* 1991;17:357-367.
- Gelman N, Ewing JR, Gorell JM, Spickler EM, Solomon EG. Interregional variation of longitudinal relaxation rates in human brain at 3.0 T: relation to estimated iron and water contents. *Magn Reson Med* 2001;45:71-79.
- Ewing JR, Aryal M, Bagher-Ebadian H, Panda S, Karki K, Nagaraja TN, Cabral G, Fenstermacher JD, Mikkelsen T. Dynamic contrast enhanced MRI at 7 T in a rat model of cerebral glioma: data analysis and model selection. In Proceedings of the 20th Annual Meeting of ISMRM, Melbourne, Australia, 2012. p. 1530.
- Sykova E, Nicholson C. Diffusion in brain extracellular space. *Physiol Rev* 2008;88:1277-1340.
- Jain R, Ellika SK, Scarpace L, Schultz LR, Rock JP, Gutierrez J, Patel SC, Ewing J, Mikkelsen T. Quantitative estimation of permeability surface-area product in astroglial brain tumors using perfusion CT and correlation with histopathologic grade. *AJNR Am J Neuroradiol* 2008;29:694-700.
- Pickup S, Chawla S, Poptani H. Quantitative estimation of dynamic contrast enhanced MRI parameters in rat brain gliomas using a dual surface coil system. *Acad Radiol* 2009;16:341-350.
- Chenevert TL, Stegman LD, Taylor JM, Robertson PL, Greenberg HS, Rehemtulla A, Ross BD. Diffusion magnetic resonance imaging: an early surrogate marker of therapeutic efficacy in brain tumors. *J Natl Cancer Inst* 2000;92:2029-2036.
- Farace P, Merigo F, Fiorini S, Nicolato E, Tambalo S, Daducci A, Degrassi A, Sbarbati A, Rubello D, Marzola P. DCE-MRI using small-molecular and albumin-binding contrast agents in experimental carcinomas with different stromal content. *Eur J Radiol* 2011;78:52-59.
- Li X, Rooney WD, Springer CS Jr. A unified magnetic resonance imaging pharmacokinetic theory: intravascular and extracellular contrast reagents. *Magn Reson Med* 2005;54:1351-1359.
- Landis CS, Li X, Telang FW, Coderre JA, Micca PL, Rooney WD, Latour LL, Vetek G, Palyka I, Springer CS Jr. Determination of the MRI contrast agent concentration time course in vivo following bolus injection: effect of equilibrium transcytolemmal water exchange. *Magn Reson Med* 2000;44:563-574.
- Landis CS, Li X, Telang FW, Molina PE, Palyka I, Vetek G, Springer CS. Equilibrium transcytolemmal water-exchange kinetics in skeletal muscle in vivo. *Magn Reson Med* 1999;42:467-478.
- Li X, Rooney WD, Varallyay CG, et al. Dynamic-contrast-enhanced-MRI with extravasating contrast reagent: rat cerebral glioma blood volume determination. *J Magn Reson* 2010;206:190-199.
- Paudyal R, Poptani H, Cai K, Zhou R, Glickson JD. Impact of transvascular and cellular-interstitial water exchange on dynamic contrast-enhanced magnetic resonance imaging estimates of blood to tissue transfer constant and blood plasma volume. *J Magn Reson Imaging* 2013;37:435-444.