

Henry Ford Health

## Henry Ford Health Scholarly Commons

---

Emergency Medicine Articles

Emergency Medicine

---

2-1-2021

### Systemic thrombolysis for refractory cardiac arrest due to presumed myocardial infarction

Joseph A. Hamera

Noah B. Bryant

Mark S. Shievitz

*Henry Ford Health*, mshievi1@hfhs.org

David A. Berger

Follow this and additional works at: [https://scholarlycommons.henryford.com/emergencymedicine\\_articles](https://scholarlycommons.henryford.com/emergencymedicine_articles)

---

#### Recommended Citation

Hamera JA, Bryant NB, Shievitz MS, and Berger DA. Systemic thrombolysis for refractory cardiac arrest due to presumed myocardial infarction. *Am J Emerg Med* 2021; 40:226.e223-226.e225.

This Article is brought to you for free and open access by the Emergency Medicine at Henry Ford Health Scholarly Commons. It has been accepted for inclusion in Emergency Medicine Articles by an authorized administrator of Henry Ford Health Scholarly Commons.



# Systemic thrombolysis for refractory cardiac arrest due to presumed myocardial infarction

Joseph A. Hamera, MD<sup>a,b</sup>, Noah B. Bryant, MD<sup>c,d</sup>, Mark S. Shievitz, MD<sup>e</sup>, David A. Berger, MD<sup>a,c,\*</sup>

<sup>a</sup> Department of Emergency Medicine, Beaumont Hospital-Royal Oak, Royal Oak, MI, United States of America

<sup>b</sup> Department of Emergency Medicine, University of Michigan Health System, Ann Arbor, MI, United States of America

<sup>c</sup> Oakland University William Beaumont School of Medicine, Rochester, MI, United States of America

<sup>d</sup> Department of Emergency Medicine, Beaumont Hospital-Grosse Pointe, Grosse Pointe, MI, United States of America

<sup>e</sup> Department of Emergency Medicine, Henry Ford Medical Center-Fairline, Dearborn, MI, United States of America

## ARTICLE INFO

### Article history:

Received 14 June 2020

Received in revised form 8 July 2020

Accepted 16 July 2020

### Keywords:

Thrombolysis

Out of hospital cardiac arrest

## ABSTRACT

The empiric usage of systemic thrombolysis for refractory out of hospital cardiac arrest (OHCA) is considered for pulmonary embolism (PE), but not for undifferentiated cardiac etiology [1, 2]. We report a case of successful resuscitation after protracted OHCA with suspected non-PE cardiac etiology, with favorable neurological outcome after empiric administration of systemic thrombolysis.

A 47-year-old male presented to the emergency department (ED) after a witnessed OHCA with no bystander cardiopulmonary resuscitation (CPR). His initial rhythm was ventricular fibrillation (VF) which had degenerated into pulseless electrical activity (PEA) by ED arrival. Fifty-seven minutes into his arrest, we gave systemic thrombolysis which obtained return of spontaneous circulation (ROSC). He was transferred to the coronary care unit (CCU) and underwent therapeutic hypothermia. On hospital day (HD) 4 he began following commands and was extubated on HD 5. Subsequent percutaneous coronary intervention (PCI) revealed non-obstructive stenosis in distal LAD. He was discharged home directly from the hospital, with one-month cerebral performance category (CPC) score of one. He was back to work three months post-arrest.

Emergency physicians (EP) should be aware of this topic since we are front-line health care professionals for OHCA. Thrombolytics have the advantage of being widely available in ED and therefore offer an option on a case-by-case basis when intra-arrest PCI and ECPR are not available. This case report adds to the existing literature on systemic thrombolysis as salvage therapy for cardiac arrest from an undifferentiated cardiac etiology. The time is now for this treatment to be reevaluated.

© 2020 Elsevier Inc. All rights reserved.

## 1. Case report

The American Heart Association (AHA) stated fibrinolysis can be considered for cardiac arrest secondary to suspected or confirmed pulmonary embolism (PE), though not for undifferentiated refractory arrest [1,2]. Optimism for thrombolysis for OHCA from suspected acute myocardial infarction (AMI) abruptly halted after 2008 double blinded placebo controlled (DBPC) randomized controlled trial (RCT) reported futility and increased intracranial hemorrhage (ICH) [3–6]. Thereafter, the authors are not aware of further RCT in this topic.

We report a case of successful resuscitation after protracted OHCA with suspected AMI etiology, with favorable neurological outcome after administration of alteplase tissue plasminogen activator (tPA).

Our objective is to explore systemic thrombolysis as a salvage therapy for refractory OHCA despite maximal conventional resuscitative efforts.

A 47-year-old male presented to the ED after a witnessed OHCA without bystander CPR. Emergency medical services (EMS) arrived within 10 min and found patient in ventricular fibrillation (VF). He was defibrillated seven times without ROSC; arrived at ED 29 min later in PEA. Advanced cardiac life support continued and he was intubated. He did have three-minute ROSC (37 min into his arrest) where ECG revealed anterolateral AMI.

After 57 min of arrest, 50 mg of intravenous tPA was administered and 11 min later he developed ROSC with sinus tachycardic rhythm, for a total of 68 min downtime. Fifteen minutes later became bradycardic and second 50 mg dose of tPA was administered.

During the remainder of ED stay he developed hypoxia and underwent right tube thoracostomy (suspected CPR-related pneumothorax). Computed tomography (CT) head and CT angiography chest was negative for ICH and PE, respectively. He was weaned off vasopressors in the ED.

\* Corresponding author at: 3601 W 13 Mile Rd., Royal Oak, MI 48073, United States of America.

E-mail address: [David.berger@beaumont.edu](mailto:David.berger@beaumont.edu) (D.A. Berger).

In CCU, he underwent therapeutic hypothermia. He remained comatose until HD four when he began to follow commands in lower extremities. The following morning, he had spontaneous eye opening and was extubated 24 h later. He transferred out of CCU on HD 12, MRI heart on HD 21 suggested myocarditis and cardiac catheterization on HD 22 found non-obstructing disease in the Left Anterior Descending artery.

He was discharged home from the hospital on HD 32 with no residual neurological deficits and cerebral performance category (CPC) score was one. He returned to work within three months post-arrest.

AMI and PE represent 65% of OHCA [7]. The most likely cause of this patient's cardiac arrest was AMI, and Otani described that in witnessed OHCA with initial shockable rhythm, greater than 50% of those are "resistant" to conventional CPR are from AMI [8]. The treatment of choice for AMI is percutaneous coronary intervention (PCI), but infrequently available during cardiac arrest.

Clinicians have reported positive findings utilizing thrombolysis for OHCA in a multitude of research designs. This includes a prospective DBPC RCT [5], before/after prospective study [3], a retrospective propensity-matched analysis [9], a large case series [10] and case reports as recent as 2015 [11–14]. A 2019 meta-analysis reported higher hospital admission rates in patients given thrombolysis, though additional outcomes for efficacy and safety raised concern [15]. The authors are uncertain whether this meta-analysis' higher "survival to admission" in thrombolytic recipients, but similar 24-h survival, could be reflective of early withdrawal of life sustaining treatment (WLST) or higher severity of illness in the thrombolytic recipients. This meta-analysis was also heavily weighted to the neg RCT in 2008, most notably for "survival to discharge" and "bleeding complications" were weighted 66.2% and 78.9%, respectively [6]. This 2008 trial also used Tenecteplase which is difficult to directly compare to prior successful work with Alteplase [3,4].

The probability of achieving a good outcome in cardiac arrest is optimized by a series of interventions starting at the time of arrest [16]. This case illustrates a potential workaround when prehospital care has preserved viability and thrombolytics seem to offer little downside [4].

One intervention which gained popularity is extracorporeal membrane oxygenation (ECMO) with CPR, collectively called ECPR. Data for ECPR for OHCA is only observational trials, and without RCT [17–19]. Our center has ECMO capability but at this time does not offer ECPR. Trials are underway to evaluate its cost-effectiveness and feasibility [20].

Also observed in this case is delayed neurological recovery after cardiac arrest, which contrasts with the fact that WLST remaining a major cause of death of comatose OHCA patients [21]. We recognize that despite our case's CPC-1 survival, no bystander-involved protracted CPR with late administration of systemic thrombolysis, the scale appeared tipped towards futility.

The authors readily acknowledge the substantial limitations of the literature on intra-arrest thrombolytics for OHCA, including publication bias. Since the 2008 RCT, other links in the "chain of survival" have been re-examined and the authors are uncertain if the optimization of other links in the chain, has any effect on utility of intra-arrest fibrinolysis [22,23]. This case report builds on the existing literature on systemic thrombolysis as salvage therapy for cardiac arrest from an undifferentiated cardiac etiology. The time is now for this treatment to be reevaluated.

## Funding disclosure

This research did not receive any specific grant funding from agencies in the public, commercial, or not-for profit sectors.

Our article has never been previously published in any form, except as abstract at the Society of Critical Care Medicine Critical Care Congress on 2/27/2018.

## Author contribution statement

All listed authors have contributed to this manuscript process as follows:

\*JAH contributed to conceptualization, original draft and review/editing.

\*NBB contributed to review/editing and clinically managed the case.

\*MSS contributed to review/editing and clinically managed the case.

\*DAB contributed to conceptualization, original draft, review/editing, clinically managed the case and supervision.

## Declaration of Competing Interest

None.

## Acknowledgements

The authors wish to acknowledge the support of our institution's therapeutic hypothermia team, including Dr. Robert Swor, Dr. Kelly Sawyer and Ryan Paternoster. We also are grateful for the bedside nurses, cardiologists, neurologists who expertly manage these cases after the patient leaves our Emergency Department. Your clinical and research excellence inspires us all.

## References

- [1] Neumar RW, et al. Part 8: adult advanced cardiovascular life support: 2010 American heart association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2010;122(18 Suppl 3):S729–67.
- [2] Lavonas EJ, et al. Part 10: special circumstances of resuscitation: 2015 American heart association guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2015;132(18 Suppl 2):S501–18.
- [3] Bottiger BW, et al. Efficacy and safety of thrombolytic therapy after initially unsuccessful cardiopulmonary resuscitation: a prospective clinical trial. *Lancet*. 2001;357(9268):1583–5.
- [4] Lederer W, et al. Recombinant tissue plasminogen activator during cardiopulmonary resuscitation in 108 patients with out-of-hospital cardiac arrest. *Resuscitation*. 2001;50(1):71–6.
- [5] Fatovich DM, Dobb GJ, Clugston RA. A pilot randomised trial of thrombolysis in cardiac arrest (the TICA trial). *Resuscitation*. 2004;61(3):309–13.
- [6] Bottiger BW, et al. Thrombolysis during resuscitation for out-of-hospital cardiac arrest. *N Engl J Med*. 2008;359(25):2651–62.
- [7] Keuper W, et al. Reperfusion therapy in out-of-hospital cardiac arrest: current insights. *Resuscitation*. 2007;73(2):189–201.
- [8] Otani T, et al. Resistance to conventional cardiopulmonary resuscitation in witnessed out-of-hospital cardiac arrest patients with shockable initial cardiac rhythm. *J Cardiol*. 2016;68(2):161–7.
- [9] Renard A, et al. Impact of fibrinolysis on immediate prognosis of patients with out-of-hospital cardiac arrest. *J Thromb Thrombolysis*. 2011;32(4):405–9.
- [10] Arntz HR, et al. Out-of-hospital thrombolysis during cardiopulmonary resuscitation in patients with high likelihood of ST-elevation myocardial infarction. *Resuscitation*. 2008;76(2):180–4.
- [11] Reyner K, et al. Successful use of intra-arrest thrombolysis for electrical storm due to acute myocardial infarction. *Am J Emerg Med*. 2015;33(7) (p. 990.e5–8).
- [12] Tiffany PA, Schultz M, Stueven H. Bolus thrombolytic infusions during CPR for patients with refractory arrest rhythms: outcome of a case series. *Ann Emerg Med*. 1998;31(1):124–6.
- [13] Duchateau FX, et al. Out-of-hospital thrombolytic therapy during cardiopulmonary resuscitation in refractory cardiac arrest due to acute myocardial infarction. *Eur J Emerg Med*. 2001;8(3):241–3.
- [14] Sheth A, et al. Bolus thrombolytic infusion during prolonged refractory cardiac arrest of undiagnosed cause. *Emerg Med J*. 2006;23(3):e19.
- [15] Wang Y, et al. Can systemic thrombolysis improve prognosis of cardiac arrest patients during cardiopulmonary resuscitation? A systematic review and meta-analysis. *J Emerg Med*. 2019;57(4):478–87.
- [16] Cummins RO, et al. Improving survival from sudden cardiac arrest: the "chain of survival" concept. A statement for health professionals from the advanced cardiac life support subcommittee and the emergency cardiac care committee, American Heart Association. *Circulation*. 1991;83(5):1832–47.
- [17] Panchal AR, et al. 2019 American Heart Association focused update on advanced cardiovascular life support: use of advanced airways, vasopressors, and extracorporeal cardiopulmonary resuscitation during cardiac arrest: an update to the American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2019;140(24):e881–94.

- [18] Stub D, et al. Refractory cardiac arrest treated with mechanical CPR, hypothermia, ECMO and early reperfusion (the CHEER trial). *Resuscitation*. 2015;86:88–94.
- [19] Bartos JA, et al. Surviving refractory out-of-hospital ventricular fibrillation cardiac arrest: critical care and extracorporeal membrane oxygenation management. *Resuscitation*. 2018;132:47–55.
- [20] Bol ME, et al. Early initiation of extracorporeal life support in refractory out-of-hospital cardiac arrest: design and rationale of the INCEPTION trial. *Am Heart J*. 2019;210:58–68.
- [21] Geocadin RG, et al. Standards for studies of neurological prognostication in comatose survivors of cardiac arrest: a scientific statement from the American Heart Association. *Circulation*. 2019;140(9):e517–42.
- [22] Bobrow BJ, et al. Chest compression–only CPR by lay rescuers and survival from out-of-hospital cardiac arrest. *JAMA*. 2010;304(13):1447–54.
- [23] Nielsen N, et al. Targeted temperature management at 33 degrees C versus 36 degrees C after cardiac arrest. *N Engl J Med*. 2013;369(23):2197–206.