Henry Ford Health [Henry Ford Health Scholarly Commons](https://scholarlycommons.henryford.com/)

[Public Health Sciences Articles](https://scholarlycommons.henryford.com/publichealthsciences_articles) [Public Health Sciences](https://scholarlycommons.henryford.com/publichealthsciences)

4-29-2021

Expression and regulatory roles of lncRNAs in G-CIMP-low vs G-CIMP-high Glioma: an in-silico analysis

Indrani Datta Henry Ford Health, idatta1@hfhs.org

Houtan Noushmehr Henry Ford Health, hnoushm1@hfhs.org

Chaya Brodie Henry Ford Health, CBRODIE1@hfhs.org

Laila M. Poisson Henry Ford Health, lpoisso1@hfhs.org

Follow this and additional works at: [https://scholarlycommons.henryford.com/](https://scholarlycommons.henryford.com/publichealthsciences_articles?utm_source=scholarlycommons.henryford.com%2Fpublichealthsciences_articles%2F262&utm_medium=PDF&utm_campaign=PDFCoverPages) [publichealthsciences_articles](https://scholarlycommons.henryford.com/publichealthsciences_articles?utm_source=scholarlycommons.henryford.com%2Fpublichealthsciences_articles%2F262&utm_medium=PDF&utm_campaign=PDFCoverPages)

Recommended Citation

Datta I, Noushmehr H, Brodie C, and Poisson LM. Expression and regulatory roles of lncRNAs in G-CIMPlow vs G-CIMP-high Glioma: an in-silico analysis. J Transl Med 2021; 19(1):182.

This Article is brought to you for free and open access by the Public Health Sciences at Henry Ford Health Scholarly Commons. It has been accepted for inclusion in Public Health Sciences Articles by an authorized administrator of Henry Ford Health Scholarly Commons.

RESEARCH

Open Access

Expression and regulatory roles of lncRNAs in G-CIMP-low vs G-CIMP-high Glioma: an in-silico analysis

Indrani Datta^{1,2}, Houtan Noushmehr², Chaya Brodie² and Laila M. Poisson^{1,2[*](http://orcid.org/0000-0002-3409-6536)}

Abstract

Background: Clinically relevant glioma subtypes, such as the glioma-CpG island methylator phenotype (G-CIMP), have been defned by epigenetics. In this study, the role of long non-coding RNAs in association with the poorprognosis G-CMIP-low phenotype and the good-prognosis G-CMIP-high phenotype was investigated. Functional associations of lncRNAs with mRNAs and miRNAs were examined to hypothesize infuencing factors of the aggressive phenotype.

Methods: RNA-seq data on 250 samples from TCGA's Pan-Glioma study, quantifed for lncRNA and mRNAs (GEN-CODE v28), were analyzed for diferential expression between G-CIMP-low and G-CIMP-high phenotypes. Functional interpretation of the diferential lncRNAs was performed by Ingenuity Pathway Analysis. Spearman rank order correlation estimates between lncRNA, miRNA, and mRNA nominated diferential lncRNA with a likely miRNA sponge function.

Results: We identified 4371 differentially expressed features (mRNA = 3705; lncRNA = 666; FDR < 5%). From these, the protein-coding gene *TP53* was identifed as an upstream regulator of diferential lncRNAs PANDAR and PVT1 $(p=0.0237)$ and enrichment was detected in the "development of carcinoma" ($p=0.0176$). Two lncRNAs (HCG11, PART1) were positively correlated with 342 mRNAs, and their correlation estimates diminish after adjusting for either of the target miRNAs: hsa-miR-490-3p, hsa-miR-129-5p. This suggests a likely sponge function for HCG11 and PART1.

Conclusions: These fndings identify diferential lncRNAs with oncogenic features that are associated with G-CIMP phenotypes. Further investigation with controlled experiments is needed to confrm the molecular relationships.

Keywords: Long non-coding RNAs, Glioma, G-CIMP subtypes

Background

Glioma, a tumor of glial cells, is the most aggressive form of tumor of the central nervous system (CNS). Historically glioma has been described by histologic features and malignancy grading. Glioblastoma (GBM) is grade IV disease, typically with necrotic regions, conferring poor overall survival (15.5% at 2 years for adult GBM, 95% CI:

*Correspondence: lpoisso1@hfhs.org

¹ Department of Public Health Sciences, Center for Bioinformatics, Henry Ford Health System, 1 Ford Place, 3C, Detroit, MI 48202, USA

15.1%–15.9%) [\[1](#page-8-0), [2\]](#page-8-1). Difuse gliomas, astrocytoma and oligodendroglioma of grade 2 or 3, are characterized by varying degrees of aggressiveness and extensive infltrative growth in the surrounding CNS parenchyma [[1,](#page-8-0) [2](#page-8-1)]. Recently, the World Health Organization (WHO) added the presence of one of the recurrent point mutations in the isocitrate dehydrogenase genes (*IDH1* or *IDH2*) and co-deletion of chromosomal arms 1p/19q to the glioma diagnosis criteria [[3\]](#page-8-2). Yet, even the refned molecular diagnosis classifcations do not fully explain the heterogeneous clinical phenotypes of these tumors.

© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit [http://creativeco](http://creativecommons.org/licenses/by/4.0/) [mmons.org/licenses/by/4.0/.](http://creativecommons.org/licenses/by/4.0/) The Creative Commons Public Domain Dedication waiver ([http://creativecommons.org/publicdomain/](http://creativecommons.org/publicdomain/zero/1.0/) [zero/1.0/\)](http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Full list of author information is available at the end of the article

With recent advances in genomics, molecular subtypes are able to be further refned. In glioma, characterization of the epigenome by DNA methylation assay has been useful in the stratifcation and integration of molecular and phenotypic features [[3\]](#page-8-2). One such sub-classifcation, known as the CpG island methylator phenotype (CIMP), is defned by genome-wide hypermethylation of CpG Islands (CGI) and was frst defned in the context of colorectal cancer $[4, 5]$ $[4, 5]$ $[4, 5]$ $[4, 5]$ $[4, 5]$. The glioma-CIMP (G-CIMP) subtype was frst described by Noushmehr et al. [\[6](#page-8-5)] in glioblastoma (GBM; Grade 4 glioma) and then in lowergrade gliomas (LGG; Grades 2, 3). Compared to G-CIMP negative tumors, several studies found that G-CIMP positive subtypes were typically associated with younger patients and with *IDH*-mutant gliomas without 1p/19q co-deletion [\[6](#page-8-5), [7](#page-8-6)]. Tis *IDH*-mutant G-CIMP positive subtype has now been further refned into two distinct subgroups, G-CIMP-low (10% of *IDH*-mutant, 1p/19q intact tumor) and G-CIMP-high (90% of *IDH*-mutant, 1p/19q intact tumors), with 'low' and 'high' designations determined by a low or high degree of DNA methylation, respectively. As opposed to the characteristically higher survival rate of IDH-mutant glioma, G-CIMP-low tumors have survival rates that are closer to that of GBM. Even though the two G-CMIP subtypes have molecularly distinct methylation patterns, factors driving the diference in prognosis are yet unknown. It is assumed that DNA methylation patterns are associated with transcriptomic patterns, including non-coding RNAs. With recent advances in RNA sequencing various RNA species can be quantifed (e.g., coding messenger RNA [mRNA], micro RNA [miRNA;22–24 bp], and long non-coding RNA [lncRNA,>200 bp]). Among the RNA species, epigenetic regulators such as long non-coding RNA (lncRNA) have gained attention in recent years in cancer research.

LncRNAs are minimally 200-nucleotide RNA, with no known translational capacity. They have drawn attention due to their potential to regulate many cellular activities, as well as gene expression, in biological and pathological processes. Acting as cellular address codes, lncRNAs transfer proteins to their appropriate chromosomal location or fold them into higher order structures as target recognition for chromatin remodeling. In glioma, lncR-NAs have been associated with oncogenesis and prognosis $[8, 9]$ $[8, 9]$ $[8, 9]$ $[8, 9]$, such as in the recent global analysis of lncRNAs in TCGA grade 2–4 gliomas that identifed a panel of 64 lncRNAs associated with prognosis [\[10\]](#page-8-9). Among the specifc lncRNAs studied, HOTAIR—a well-known, highlyexpressed lncRNA in breast cancer [[11\]](#page-8-10)—has been associated with biogenesis and diferentiation of gliomas [[10\]](#page-8-9). TALNEC2, a lncRNA highly expressed in GBMs and with poor prognosis when silenced, inhibited cell proliferation and arrested the cells in the G1\S phase of the cell cycle in patient-derived glioma cell lines [[12\]](#page-8-11). In addition, some newly discovered lncRNAs such as lncRNA ASLNC22381 and KIAA0495 [\[9](#page-8-8)] have been found in glioma tissue and cell lines.

Since each of the glioma subtypes is clinically distinct, understanding the role of associated epigenetic regulators could help to better distinguish between the groups. Diferential epigenetic regulators may also identify biological diferences underlying the phenotypes or suggest novel therapeutic targets. In this study we aimed to identify diferentially expressed lncRNA between G-CIMPhigh and G-CIMP-low glioma, using RNA sequencing data from the glioblastoma (GBM; Grade 4 glioma) and lower grade glioma (LGG; Grades 2 and 3) arms of The Cancer Genome Atlas (TCGA). As the functional roles of most lncRNAs are poorly understood, we evaluated lncRNA involvement by gene-set enrichment from an available functional knowledgebase. Finally, we integrated lncRNA, miRNA, and mRNA expression through correlation estimates to identify lncRNAs that may be afecting transcription level changes in relationship to miRNA by acting as a miRNA sponge.

Results

Sample details

A summary of the clinical and demographic data of the TCGA G-CMIP cohort is represented in Table [1](#page-3-0). These cases are *IDH* mutant tumors, without 1p/19q co-deletion, by defnition of the G-CMIP phenotype. As such, the majority of the cases were under 40 years old at diagnosis; most tumors had an astrocytomatous histology. Primary diagnosis for these tumors used the 2007 WHO diagnosis guidelines. The WHO 2016 diagnosis was inferred from WHO grading $(2/3)$ = Astrocytoma; 4=GBM) and molecular features (IDH mutant, without 1p/19q co-deletion).

Association of lncRNA expression with G‑CIMP group

The comparison of RNA sequencing reads between glioma subtypes G-CIMP-high and G-CIMP-low identifed 4371 diferentially expressed (DE) features $(mRNA=3705, IncRNA=666)$ at a false discovery rate of 0.05. Figure [1](#page-3-1) shows a heatmap of the DE lncRNAs between GCIMP-high and GCIMP-low groups. Here the expression level is standardized per row with yellow high and blue low. The rows are ordered by hierarchical clustering of lncRNA expression. The majority of the 666 diferential lncRNAs identifed are highly expressed, with a maximum fold change of expression in G-CIMP-low tumors four times that of G-CIMP-high tumors.

The biological role of the DE lncRNA was explored via pathway analyses. Of the 666 DE lncRNAs, 44 were identifed by lncRNA gene symbols in the QIAGEN

	G-CIMP High	G-CIMP Low
Number of Cases	234	16
Age		
Under 40 (%)	149 (63.7%)	10 (62.5%)
Over 40 (%)	80 (34.2%)	6(37.5%)
Unknown (%)	5(2.1%)	$0(0.0\%)$
Gender		
Male (%)	133 (56.8%)	7 (43.8%)
Female (%)	100 (42.7%)	$9(56.2\%)$
Unknown (%)	$1(0.4\%)$	$0(0.0\%)$
WHO 2007 Histology		
Oligodendroglioma	39 (16.7%)	$0(0.0\%)$
Oligoastrocytoma	74 (31.6%)	$1(6.2\%)$
Astrocytoma	118 (50.4%)	10 (62.5%)
Glioblastoma	$2(0.9\%)$	5(31.2%)
WHO Grade		
\mathfrak{D}	111 (47.4%)	
3	92 (39.3%)	8 (50.0%)
$\overline{4}$	$2(0.9\%)$	5(31.2%)
WHO 2016 Diagnosis		
IDHmut - Astroctyoma	232 (99.1%)	11 (68.8%)
IDHmut - Glioblastoma	$2(0.9\%)$	5 (31.2%)

Table 1 Patient characteristics for the 250 primary glioma diagnoses in this study

Knowledge Base and therefore available for analysis by the Ingenuity Pathway Analysis tools. Set-enrichment analyses identifed protein-coding gene *TP53* as an upstream regulator of DE lncRNA PANDAR and PVT1 $(p=0.0237;$ Fig. [2a](#page-4-0)). In addition, "development of carcinoma" was identifed as an enriched disease category (Fig. [2b](#page-4-0)). Specifcally, four oncology-related sets were among the most enriched disease and biological function categories (Table [2\)](#page-5-0): 'breast or colorectal cancer' (9 DE lncRNAs, $p=0.0049$), 'development of digestive organ tumor' (8 DE lncRNAs, $p=0.0122$), 'development of carcinoma' (9 DE lncRNAs, $p=0.0176$), and 'malignant genitourinary solid tumor' (8 DE lncRNAs, $p = 0.0254$). A top constructed biological network associated with the DE lncRNA list was associated with cell death and survival, cellular growth, and proliferation cellular development. This network was based on 6 DE lncRNAs with 29 genes from the IPA knowledgebase (enrichment score of 14; Fig. [2](#page-4-0)c). Network scores are based on the networkeligible molecules in the analysis. Scores are inversely related to the probability of fnding the selected networkeligible molecules in a given network by random chance.

Nomination of lncRNA as a miRNA sponge

Since less than 10% of the lncRNAs had known function in the pathway analysis we also used a data driven

approach to identify lncRNA functioning as miRNA sponges. To identify lncRNA:miRNA:mRNA triplets (sponge relationships), Spearman rank order correlation $(r_{x,y})$ was estimated on expression levels between each mRNA and DE lncRNA. As described in the methods, the $r_{x,y}$ > 0.5 threshold resulted in 580 (lncR-NAs) and 14,425 (mRNAs) selected. After fltering the correlated pairs to only those with a common miRNA target for the lncRNA and mRNA, 121,276 triplets were constructed from 15 lncRNAs, 6777 mRNAs, and 201 miRNAs. To assess if a triplet was likely to refect a sponge relationship, the partial correlation $(r_{x,v|z})$ between each lncRNA and the correlated mRNA was estimated, controlling for the predicted common miRNA. The distribution of the influence of miRNA on the lncRNA:mRNA correlation, specifcally $Sz = r_{x,y} - r_{x,y|z}$, is plotted in Additional file [1:](#page-8-12) Fig S1. Two miRNAs fell into the 99th percentile of this Sz distribution: hsa-miR-129-5p, hsa-miR-490-3p. Associated with these two miRNAs were two lncR-NAs (HCG11, PART1), which were correlated with 290 (HCG11) and 114 (PART1) mRNAs, respectively, forming miRNA:lncRNA:mRNA trios. Thus, HCG11 and

Disease and Biofunctions	P-values	Molecules	
Breast or colorectal cancer	0.0049	C10orf25, CASC2, CRNDE, FOXD2-AS1, HCG11, HOTAIRM1, LINC00346, NORAD, PANDAR	
Development of digestive organ tumor	0.0122	C10orf25, CASC2, CRNDE, EPB41L4A-AS1, FOXD2-AS1, HOTAIRM1, PCA3, PVT1	
Development of carcinoma	0.0176	C10orf25, CASC2, CRNDE, CYTOR, EPB41L4A-AS1, FOXD2-AS1, HOTAIRM1, PCA3, PVT1	
Colorectal carcinoma	0.0231	C10orf25, CRNDE, FOXD2-AS1, HOTAIRM1	
Malignant genitourinary solid tumor	0.0254	CASC15, CASC2, CRNDE, HCG11, LINC00346, NORAD, PANDAR, PCA3	

Table 2 Biological functions from IPA knowledgebase that are enriched with DE lncRNAs

The p-values are calculated with Fisher exact test. Molecules listed here are those DE lncRNAs identifed within the functional group being assessed

PART1 were nominated as sponges for hsa-miR-129-5p and hsa-miR-490-3p, blocking their interaction with 342 unique mRNA; Fig. [3](#page-5-1).

Discussion

In our investigation of the non-coding transcriptome between G-CIMP-high and G-CMIP-low glioma subclasses, 666 diferential lncRNAs were identifed; some of these lncRNA had previously been associated with oncogenic activities in cancer. The expression levels of the majority of these lncRNAs were lower in G-CIMPhigh tumors. The up-regulation in G-CIMP-low tumors may relate to oncogenic activities leading to their aggressive phenotype and poor survival. A previously known relationship between TP53 (p53) and PVT1, as a p53-induced target gene, was observed by examination of upstream regulators of the diferential lncRNA [[13\]](#page-8-13). Further, we nominated two lncRNAs as having potential sponge activity for two microRNAs.

TP53 as a regulator of PANDAR and PVT1

TP53 is a well-known gene for a tumor suppressor protein p53 that participates in diverse cellular functions including cell cycle arrest, apoptosis, senescence, DNA repair, and changes in metabolism. Mutation of TP53 is associated with a variety of human cancers including gliomas, and is found in 94 percent of IDH mutant, 1p/19q non-codeleted, glioma [[14](#page-8-14)]. High expression of PVT1, a long non-coding RNA located at chromosome 8a24.21, has been associated with several mutations of TP53 in diffuse glioma $[8]$ $[8]$. In these data, the expression of TP53 and PVT1 are positively correlated, with highest expression of each occurring in GCIMP-low, see Additional fle [2:](#page-8-15) Fig S2. Many studies have shown evidence of carcinogenic activity of PVT1 in various cancers, such as negatively modulating miRNA by acting as a competing endogenous RNA or acting as a sponge to promote tumor efects [[15](#page-8-16)]. Of importance to glioma, PVT1 has been implicated in regulating levels the proto-oncogene MYC to promote tumorigenesis $[16]$ $[16]$ $[16]$. The role of MYC in glioma has been well established, both in vivo and in vitro, such that MYC inhibition suppresses glioma formation, restricts glioma cell proliferation and improves survival [[17\]](#page-8-18). The relationship between TP53 and PANDAR is less understood, though human p53 [TP53] protein is necessary for expression of human PANDA [PANDAR] lncRNA. PANDAR is a promoter of CDKN1A antisense DNA damage activated RNA and increased expression of PANDAR has been indicated to predict poor prognosis in cervical and gastric cancer $[18, 19]$ $[18, 19]$ $[18, 19]$ $[18, 19]$ $[18, 19]$. Recently, a study published showed CDKN2A, a gene which belongs to same family as CDKN1A, often deleted in G-CIMP-low tumors as compared to G-CIMP-high [[20\]](#page-8-21). While the role of PANDAR has been evaluated in many cancers [[21](#page-8-22)] its association with glioma has not been studied.

Network focused on cell growth, proliferation, survival and death

Cell death is one of the primary mechanisms studied in cancer as disruption of this process can facilitate tumorigenesis, promote proliferation, and lead to resistance to anticancer therapy. One of the top gene–gene networks

generated from the IPA knowledgebase was associated with the biological processes of cell death, survival, cell growth and proliferation (Fig. $2c$ $2c$). This network was constructed from 6 DE lncRNAs (PVT1, PANDAR, FOXD2- AS1, CYTOR CRNDE, HOTAIRM1) and 29 mRNAs that interact directly (solid lines) or indirectly via an intermediate gene (dotted line). Among the interacting mRNAs, the NOTCH1 gene's role in glioma pathogenesis is well established as it afects glioma tumorigenesis and maintenance. Several studies during recent years reported dysregulated NOTCH signaling activity (NOTCH 1–4) in human brain tumors [[22\]](#page-8-23). In an extensive study on the functional role of NOTCH1 in gliomas, it is observed that NOTCH1 is involved in maintaining glioma cells in an undiferentiated state, and its inhibition leads to cells maturing into a less aggressive phenotype [\[22](#page-8-23)]. Also, in the network (Fig. [2c](#page-4-0)) is MYC, a proto-oncogene that encodes nuclear phosphoprotein control as a transcription factor for its target genes. As described above, we see again the association between MYC and the PAN-DAR lncRNA.

Enrichment of oncogenic function

The biofunction "development of carcinoma" was one of the top cancer-related enriched biofunctions (p-value of 0.0176), with 8 DE lncRNAs: PVT1, CASC2, PCA3, EPB41L4A-AS1, C10orf25, CYTOR, FOXD2-AS1, and CRNDE. The increased expression of PVT1, CYTOR, FOXD2-AS1, and CRNDE were seen in various cancers, similarly these lncRNAs were all up-regulated in G-CIMP-low suggesting their more oncogenic activity leads to poor prognosis compared to G-CIMP-high. While decreased expression has been seen in CASC2 and PCA3 in cancer, these lncRNAs were down-regulated in G-CIMP-low suggesting their tumor suppressor potential [[23–](#page-8-24)[27](#page-9-1)]. Table [2](#page-5-0) shows other carcinogenic functions from the enrichment of DE lncRNAs with IPA knowledgebase.

Nominated lncRNA sponge activity

Research in many cancers has shown that lncRNAs can regulate mRNA expression levels indirectly through miRNA, by acting as a miRNA sponge. LncRNAs HCG11 and PART-1 were identifed as potential sponges for the miRNAs hsa-miR-490-3p and hsa-miR-129-5p (Fig. [3](#page-5-1)). Previous studies [\[28](#page-9-2)] have shown lncRNA HCG11 to be down-regulated in glioma tissues and cells, and this was associated with a lower survival rate in glioma patients. The observed mechanism is for lncRNA HCG11 to suppress growth of glioma was by acting with the miR-4425 to release MTA3. MiR-4425 is up-regulated in glioma tissues and a high expression of miR-4425 is associated with an unfavorable prognosis in glioma [[28\]](#page-9-2). Here we also see decreased expression of HCG11 in our lower survival G-CIMP-low group and propose an interaction with miR-490 and miR-129. LncRNA PART-1 has been shown to have oncogenic activity in colorectal cancer [\[29](#page-9-3)], but was identifed as positively associated with GBM prognosis [[30](#page-9-4)], such that decreased PART-1 predicted decreased survival time [\[31](#page-9-5)]. In this study we observed that PART-1 expression was lower in the poor-prognosis G-CIMP-low tumors, compared to G-CIMP-high tumors, which aligns with the observed relationship in GBM. Neither lncRNA HCG11 nor PART1 have been investigated for a relationship with hsa-miR-129-5p and hsa-miR-490-3p in glioma. From prior research in lung and hepatocellular carcinoma, increased hsa-miR-490-3p has been implicated in cell migration and cancer progression to metastatic disease [\[32](#page-9-6), [33](#page-9-7)]. Reduction of hsa-miR-490-3p through sponge action of lncRNAs suggests a more favorable outcome, which we see in G-CMIP-high. In contrast, hsamiR-129-5p has been shown to inhibit the cell cycle and induce apoptosis in glioma cell lines through inhibition of NOTCH1 and mTOR signaling [\[34](#page-9-8)]. Reduction of has-miR-129-5p through sponge action may thus allow increased proliferation, which is counter to expectation but dependent upon signaling pathways, so more study is needed.

Conclusion

This in-silico study explores the potential influence of non-coding RNA on the phenotypic diference between the G-CMIP-high and G-CMIP-low subtypes of glioma. The G-CMIP-low subtype is rare in primary glioma diagnosis, with less than 5% of all difuse glioma diagnoses identifed as G-CMIP-low. However, prior work shows that G-CMIP-high tumors may evolve to a G-CMIP-low form as the disease progresses [\[36](#page-9-9)]. With this in-silico study, we identifed 666 lncRNAs that showed a diference in mean expression between the two G-CMIP subtypes. With the IPA knowledgebase, we were able to propose the functional role of a subset of diferential lncRNAs related to progression to aggressive G-CIMPlow gliomas. In addition, we identifed an upstream regulator, TP53, a well-known tumor suppressor gene which can regulate two of the diferential lncRNAs. Unfortunately, our study is not without limitation, as only 44 of 666 lncRNAs had biological function information in the IPA knowledgebase. We were heartened that these 44 showed oncogenic relationships with genes known to have a role in glioma, however, we realize that there is much to be discovered among 622 lncRNAs with no information in IPA. Beyond IPA, we also identifed two lncRNAs as potentially having miR-sponge activity, HCG11 and PART-1. Each has prior evidence of an efect on glioma prognosis, which increases our enthusiasm for further study.

Methods

Ethics statement

RNA-sequencing datafles (TCGA Glioblastoma (GBMs) and Lower-grade glioma (LGGs)) were obtained from the Genomic Data Commons with appropriate approval from $dbGAP$ (#1904). They were acquired with a protocol approved by the Henry Ford Health System institutional review board (protocol #8718). The need for consent was waived in this secondary data analysis since primary identifers were not provided by dbGAP.

Quantifcation of mRNA & lncRNAs

Aligned sequencing reads (BAM fles) for TCGA GBMs and LGGs, generated from the Illumina Hiseq platform, were obtained from the Genomic Data Commons database (GDC) (March 2017). These Illumina raw reads had been processed through the RNA-Seq standardized pipeline at GDC. Briefy, the GDC pipeline frst converted to fastq with 'Biobambam' and then re-aligned to the GRCh38 reference genome per alignment guidelines from International Cancer Genome Consortium (ICGC), using the STAR aligner. A two-pass method was used for alignment; frst, splice-junctions were aligned separately in each read group, then the read groups were merged to obtain the fnal alignment in BAM format. Upon downloading these aligned BAM fles, we quantifed the read counts for lncRNAs and protein-coding messenger RNA (mRNA) against the reference annotation from GEN-CODE v28 $[37]$ $[37]$. This quantification was executed with the 'featurecount' function, from the R Bioconductor package 'Rsubread,' [[38\]](#page-9-11) which assigns mapped sequencing reads to genomic features. Two-hundred ffty samples (see Table [1\)](#page-3-0) and 37,281 features (22,583 mRNAs and 5729 antisense, 7845 lincRNA, 939 sense-intronic, 185 sense-overlapping lncRNA) were carried forward for further analysis.

Identifying diferentially expressed sets of mRNA & lncRNAs specifc to subtypes

The quantified expression matrix (in terms of read counts) for each sample was further fltered for low count, based on counts per million (CPM). Features with sum of expression values below the condition cutoff $(CPM < 1)$ across conditions were removed. Normalization between cases was performed on the weighted trimmed mean of the log2 expression ratios (TMM; trimmed means of M-values) using the R Bioconductor package NOISeqBio [\[39](#page-9-12)]. This normalization method assumes that the majority of the RNA features are not diferential. After pre-processing, 24,178 features were retained for analysis. The NOISeqBIO package was used to identify diferential expression per feature (mRNA and lncRNA) between G-CIMP-high and G-CIMP-low, considering the log2-ratio of the two conditions (M-value) and the value of the diference between conditions (D-value). A feature was identifed as diferential between conditions if its corresponding M and D values are likely to be higher than the expected noise, where the noise distribution is obtained from comparing all sample pairs within a condition. lncRNA were identifed as diferential between the two groups if the false discovery rate (FDR) was less than or equal to 5%. The differential mRNA and lncRNA identifed in this analysis are presented in Additional fles [3](#page-8-25) and [4.](#page-8-26)

Pathway analysis of diferential lncRNA

LncRNAs that were diferentially expressed between G-CIMP-high and G-CIMP-low tumors were further evaluated for biological functional interpretation with Ingenuity's IPA software knowledgebase $[40]$ $[40]$. The core enrichment analysis was performed using all lncRNAs from the Gencodev28 annotation, described above, as the reference set for the Fisher's exact test used to calculate enrichment p-values.

Integrative analysis of lncRNAs, miRNAs and mRNA to predict miRNA sponge activity

In scenarios where a lncRNA is acting as miRNA sponge, it is expected that the correlation between lncRNA and mRNA expression will be positive. Further, the correlation between lncRNA and mRNA will be dependent on miRNA expression, such that it lessens when the miRNA expression is considered. For this analysis, TCGA miRNA-seq data were downloaded from the Broad Firehose for 239 samples (G-CIMP-high=228, G-CIMP-low=11). Data had been aligned and quantifed by Broad, reporting log2 reads per million (RPM) for 2588 miRNAs. Transcriptome-wide microRNA target prediction for each lncRNA and mRNA observed to be diferentially expressed between G-CIMP-high and G-CIMP-low was obtained from MiRcode [[41\]](#page-9-14) annotation. Correlated lncRNA and mRNA pairs with a common miRNA target were retained for further analysis. To assess whether a sponge relationship is likely within each lncRNA:mRNA:miRNA trio, correlation between the lncRNA and mRNA expression was estimated, alone $(r_{x,y};$ Spearman rank order correlation) and controlling for the target miRNA expression $(r_{x,y|z};$ Spearman rank order partial correlation) $[42]$ $[42]$ $[42]$. The unconditional correlation was filtered to $r_{x,y}$ = 0.5, with p-value < = 0.05. Then a lncRNA was nominated as having a sponge function in the lncRNA:miRNA:mRNA trio if $Sz = r_{x,y} - r_{x,y|z}$ was high; here we use $Sz > 0.2$ [[42\]](#page-9-15).

Supplementary Information

The online version contains supplementary material available at [https://doi.](https://doi.org/10.1186/s12967-021-02844-z) [org/10.1186/s12967-021-02844-z.](https://doi.org/10.1186/s12967-021-02844-z)

Additional fle 1: Fig. S1. The distribution of the infuence of miRNA on the lncRNA:mRNA correlation, $Sz = r_{x,y} - r_{x,y|z}$, is plotted. For RNA triplets a reduction of correlation (S_7) of 0.2 or great were retained, in this study 0.2 is approximately the 99th percentile of the distribution of the S_z distribution.

Additional fle 2: Fig S2. Scatterplot of the expression of TP53 and PVT1 in G-CIMP-high and G-CIMP-low tumors.

Additional file 3: Table S1. Differential LncRNAs.

Additional fle 4: Table S2. Diferential_mRNAs.

Acknowledgements

Not applicable here.

Authors' contributions

The formal analysis, visualization, and preparation of the original draft was conducted by ID. Supervision was provided by LMP. All authors contributed to conceptualization and review/editing of the original draft. All authors read and approved the fnal manuscript.

Funding

This work was supported by the Henry Ford Health System, Department of Neurosurgery and the Hermelin Brain Tumor Center Foundation. Additionally, ID, LMP and HN are supported by National Institutes of Health (R01CA222146) and HN and LMP are supported by Department of Defense (CA170278).

Availability of data and materials

RNA-sequencing datafles (TCGA Glioblastoma (GBMs) and Lower-grade glioma (LGGs)) were obtained from the Genomic Data Commons with appropriate approval from dbGAP (#1904). Results datasets supporting the conclusions of this article are included in the additional fles.

Declarations

Ethics approval and consent to participate

Data were acquired from the Genomic Data Commons under a protocol approved by the Henry Ford Health System institutional review board (protocol #8718).

Consent for publication

Not applicable here.

Competing interests

The authors declare no competing interests.

Author details

¹ Department of Public Health Sciences, Center for Bioinformatics, Henry Ford Health System, 1 Ford Place, 3C, Detroit, MI 48202, USA. ² Department of Neurosurgery, Hermelin Brain Tumor Center, Henry Ford Cancer Institute, Henry Ford Health System, Detroit, USA.

Received: 11 February 2021 Accepted: 18 April 2021 Published online: 29 April 2021

References

- 1. Ostrom QT, Gittleman H, Truitt G, Boscia A, Kruchko C, Barnholtz-Sloan JS. CBTRUS Statistical Report: Primary Brain and Other Central Nervous System Tumors Diagnosed in the United States in 2011–2015. Neurooncology. 2018;20(4):1-86.
- 2. Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, et al. The 2016 World Health Organization classifcation of

tumors of the central nervous system: a summary. Acta Neuropathol. 2016;131(6):803–20.

- 3. Ceccarelli M, Barthel FP, Malta TM, Sabedot TS, Salama SR, Murray BA, et al. Molecular profling reveals biologically discrete subsets and pathways of progression in difuse glioma. Cell. 2016;164(3):550–63.
- 4. Toyota M, Ahuja N, Ohe-Toyota M, Herman JG, Baylin SB, Issa JP. CpG island methylator phenotype in colorectal cancer. Proc Natl Acad Sci USA. 1999;96(15):8681–6.
- 5. Weisenberger DJ. Characterizing DNA methylation alterations from The Cancer Genome Atlas. J Clin Investig. 2014;124(1):17–23.
- 6. Noushmehr H, Weisenberger DJ, Diefes K, Phillips HS, Pujara K, Berman BP, et al. Identifcation of a CpG island methylator phenotype that defnes a distinct subgroup of glioma. Cancer Cell. 2010;17(5):510–22.
- 7. Turcan S, Rohle D, Goenka A, Walsh LA, Fang F, Yilmaz E, et al. IDH1 mutation is sufficient to establish the glioma hypermethylator phenotype. Nature. 2012;483(7390):479–83.
- 8. Zou H, Wu LX, Yang Y, Li S, Mei Y, Liu YB, et al. lncRNAs PVT1 and HAR1A are prognosis biomarkers and indicate therapy outcome for difuse glioma patients. Oncotarget. 2017;8(45):78767–80.
- 9. Yan Y, Xu Z, Li Z, Sun L, Gong Z. An insight into the increasing role of LncRNAs in the pathogenesis of gliomas. Front Mol Neurosci. 2017;10:53.
- 10. Reon BJ, Anaya J, Zhang Y, Mandell J, Purow B, Abounader R, et al. Expression of lncRNAs in low-grade gliomas and glioblastoma multiforme: an in silico analysis. PLoS Med. 2016;13(12):e1002192.
- 11. Van Grembergen O, Bizet M, de Bony EJ, Calonne E, Putmans P, Brohee S, et al. Portraying breast cancers with long noncoding RNAs. Sci Adv. 2016;2(9):e1600220.
- 12. Brodie S, Lee HK, Jiang W, Cazacu S, Xiang C, Poisson LM, et al. The novel long non-coding RNA TALNEC2, regulates tumor cell growth and the stemness and radiation response of glioma stem cells. Oncotarget. 2017;8(19):31785–801.
- 13. Barsotti AM, Beckerman R, Laptenko O, Huppi K, Caplen NJ, Prives C. p53-Dependent induction of PVT1 and miR-1204. J Biol Chem. 2012;287(4):2509–19.
- 14. Brat DJ, Verhaak RG, Aldape KD, Yung WK, Salama SR, Cooper LA, et al. Comprehensive, integrative genomic analysis of difuse lower-grade gliomas. N Engl J Med. 2015;372(26):2481–98.
- 15. Pan X, Zheng G, Gao C. LncRNA PVT1: a Novel Therapeutic Target for Cancers. Clin Lab. 2018;64(5):655–62.
- 16. Jin K, Wang S, Zhang Y, Xia M, Mo Y, Li X, et al. Long non-coding RNA PVT1 interacts with MYC and its downstream molecules to synergistically promote tumorigenesis. Cellular and molecular life sciences: CMLS; 2019.
- 17. Annibali D, Whitfeld JR, Favuzzi E, Jauset T, Serrano E, Cuartas I, et al. Myc inhibition is effective against glioma and reveals a role for Myc in profcient mitosis. Nat Commun. 2014;5:4632.
- 18. Huang HW, Xie H, Ma X, Zhao F, Gao Y. Upregulation of LncRNA PANDAR predicts poor prognosis and promotes cell proliferation in cervical cancer. Eur Rev Med Pharmacol Sci. 2017;21(20):4529–35.
- 19. Ma P, Xu T, Huang M, Shu Y. Increased expression of LncRNA PANDAR predicts a poor prognosis in gastric cancer. Biomed Pharmacother. 2016;78:172–6.
- 20. Li KK-W, Shi Z-f, Malta TM, Chan AK-Y, Cheng S, Kwan JSH, et al. Identifcation of subsets of IDH-mutant glioblastomas with distinct epigenetic and copy number alterations and stratifed clinical risks. Neuro-Oncol Adv. 2019.
- 21. Zou Y, Zhong Y, Wu J, Xiao H, Zhang X, Liao X, et al. Long non-coding PANDAR as a novel biomarker in human cancer: a systematic review. Cell Proliferation. 2018;51:1.
- 22. Stockhausen MT, Kristoffersen K, Poulsen HS. The functional role of Notch signaling in human gliomas. Neuro Oncol. 2010;12(2):199–211.
- 23. Baldinu P, Cossu A, Manca A, Satta MP, Sini MC, Rozzo C, et al. Identifcation of a novel candidate gene, CASC2, in a region of common allelic loss at chromosome 10q26 in human endometrial cancer. Hum Mutat. 2004;23(4):318–26.
- 24. Gao J, Cao R, Mu H. Long non-coding RNA UCA1 may be a novel diagnostic and predictive biomarker in plasma for early gastric cancer. Int J Clin Exp Pathol. 2015;8(10):12936–42.
- 25. Zhang PP, Wang YQ, Weng WW, Nie W, Wu Y, Deng Y, et al. Linc00152 promotes cancer cell proliferation and invasion and predicts poor prognosis in lung adenocarcinoma. J Cancer. 2017;8(11):2042–50.
- 26. Yang X, Duan B, Zhou X. Long non-coding RNA FOXD2-AS1 functions as a tumor promoter in colorectal cancer by regulating EMT and Notch signaling pathway. Eur Rev Med Pharmacol Sci. 2017;21(16):3586–91.
- 27. Jiang H, Wang Y, Ai M, Wang H, Duan Z, Wang H, et al. Long noncoding RNA CRNDE stabilized by hnRNPUL2 accelerates cell proliferation and migration in colorectal carcinoma via activating Ras/MAPK signaling pathways. Cell Death Dis. 2017;8(6):e2862.
- 28. Zhang L, Cao Y, Kou X, Che L, Zhou X, Chen G, et al. Long non-coding RNA HCG11 suppresses the growth of glioma by cooperating with the miR-4425/MTA3 axis. J Gene Med. 2019;21(4):e3074.
- 29. Hu Y, Ma Z, He Y, Liu W, Su Y, Tang Z. PART-1 functions as a competitive endogenous RNA for promoting tumor progression by sponging miR-143 in colorectal cancer. Biochem Biophys Res Commun. 2017;490(2):317–23.
- 30. Zeng T, Li L, Zhou Y, Gao L. Exploring long noncoding RNAs in glioblastoma: regulatory mechanisms and clinical potentials. Int J Genom. 2018;2018:2895958.
- 31. Zhang XQ, Sun S, Lam KF, Kiang KM, Pu JK, Ho AS, et al. A long noncoding RNA signature in glioblastoma multiforme predicts survival. Neurobiol Dis. 2013;58:123–31.
- 32. Li J, Feng Q, Wei X, Yu Y. MicroRNA-490 regulates lung cancer metastasis by targeting poly r(C)-binding protein 1. Tumour Biol. 2016;37(11):15221–8.
- 33. Xiong L, Zhen S, Yu Q, Gong Z. HCV-E2 inhibits hepatocellular carcinoma metastasis by stimulating mast cells to secrete exosomal shuttle microR-NAs. Oncol Lett. 2017;14(2):2141–6.
- 34. Chen X, Zhang Y, Shi Y, Lian H, Tu H, Han S, et al. MiR-129 triggers autophagic fux by regulating a novel Notch-1/ E2F7/Beclin-1 axis to impair the viability of human malignant glioma cells. Oncotarget. 2016;7(8):9222–35.
- 35. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Res. 2003;13(11):2498–504.
- 36. de Souza CF, Sabedot TS, Malta TM, Stetson L, Morozova O, Sokolov A, et al. A Distinct DNA methylation shift in a subset of glioma CpG island methylator phenotypes during tumor recurrence. Cell Rep. 2018;23(2):637–51.
- 37. Frankish A, Diekhans M, Ferreira AM, Johnson R, Jungreis I, Loveland J, et al. GENCODE reference annotation for the human and mouse genomes. Nucleic Acids Res. 2019;47(D1):D766–73.
- 38. Liao Y, Smyth GK, Shi W. The R package Rsubread is easier, faster, cheaper and better for alignment and quantifcation of RNA sequencing reads. Nucleic Acids Res. 2019;47(8):e47.
- 39. Tarazona S, Furio-Tari P, Turra D, Pietro AD, Nueda MJ, Ferrer A, et al. Data quality aware analysis of diferential expression in RNA-seq with NOISeq R/Bioc package. Nucleic Acids Res. 2015;43(21):e140.
- 40. Kramer A, Green J, Pollard J Jr, Tugendreich S. Causal analysis approaches in Ingenuity Pathway Analysis. Bioinformatics (Oxford, England). 2014;30(4):523–30.
- 41. Jeggari A, Marks DS, Larsson E. miRcode: a map of putative microRNA target sites in the long non-coding transcriptome. Bioinformatics (Oxford, England). 2012;28(15):2062–3.
- 42. Olgun G, Sahin O, Tastan O. Discovering lncRNA mediated sponge interactions in breast cancer molecular subtypes. BMC Genomics. 2018;19(1):650.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional afliations.

Ready to submit your research? Choose BMC and benefit from:

- **•** fast, convenient online submission
- **•** thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- **•** gold Open Access which fosters wider collaboration and increased citations
- **•** maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

