

Henry Ford Health System

Henry Ford Health System Scholarly Commons

Neurology Articles

Neurology

7-28-2021

A Multi-center Genome-wide Association Study of Cervical Dystonia

Yan V. Sun

Chengchen Li

Qin Hui

Yunfeng Huang

Richard Barbano

See next page for additional authors

Follow this and additional works at: https://scholarlycommons.henryford.com/neurology_articles

Authors

Yan V. Sun, Chengchen Li, Qin Hui, Yunfeng Huang, Richard Barbano, Ramon Rodriguez, Irene A. Malaty, Stephen Reich, Kimberly Bambarger, Katie Holmes, Joseph Jankovic, Neepa Patel, Emmanuel Roze, Marie Vidailhet, Brian D. Berman, Mark S. LeDoux, Alberto J. Espay, Pinky Agarwal, Sarah Pirio-Richardson, Samuel A. Frank, William G. Ondo, Rachel Saunders-Pullman, Sylvain Chouinard, Stover Natividad, Alfredo Berardelli, Alexander Y. Pantelyat, Allison Brashear, Susan H. Fox, Meike Kasten, Ulrike M. Krämer, Miriam Neis, Tobias Bäumer, Sebastian Loens, Max Borsche, Simone Zittel, Antonia Maurer, Mathias Gelderblom, Jens Volkmann, Thorsten Odorfer, Andrea A. Kühn, Friederike Borngräber, Inke R. König, Carlos Cruchaga, Adam C. Cotton, Gamze Kilic-Berkmen, Alan Freeman, Stewart A. Factor, Laura Scorr, J. Douglas Bremner, Viola Vaccarino, Arshed A. Quyyumi, Christine Klein, Joel S. Perlmutter, Katja Lohmann, and Hyder A. Jinnah

RESEARCH ARTICLE

A Multi-center Genome-wide Association Study of Cervical Dystonia

Yan V. Sun, PhD,^{1,2*} Chengchen Li, MPH,¹ Qin Hui, PhD,¹ Yunfeng Huang, PhD,¹ Richard Barbano, MD, PhD,³ Ramon Rodriguez, MD,⁴ Irene A. Malaty, MD,⁵ Stephen Reich, MD,⁶ Kimberly Bambarger, BS,⁶ Katie Holmes, MPH,⁶ Joseph Jankovic, MD,⁷ Neepa J. Patel, MD,⁸ Emmanuel Roze, MD, PhD,⁹ Marie Vidailhet, MD,⁹ Brian D. Berman, MD, MS,¹⁰ Mark S. LeDoux, MD, PhD,¹¹ Alberto J. Espay, MD, MS,¹² Pinky Agarwal, MD,¹³ Sarah Pirio-Richardson, MD,¹⁴ Samuel A. Frank, MD,¹⁵ William G. Ondo, MD,¹⁶ Rachel Saunders-Pullman, MD, MPH,¹⁷ Sylvain Chouinard, MD,¹⁸ Stover Natividad, MD,¹⁹ Alfredo Berardelli, MD,²⁰ Alexander Y. Pantelyat, MD,²¹ Allison Brashear, MD, MBA,²² Susan H. Fox, PhD,²³ Meike Kasten, MD,^{24,25} Ulrike M. Krämer, PhD,²⁶ Miriam Neis, MSc,^{26,27} Tobias Bäumer, MD,^{24,28} Sebastian Loens, MD,^{24,28} Max Borsche, MD,^{24,26} Simone Zittel, MD,²⁹ Antonia Maurer,²⁹ Mathias Gelderblom, MD,²⁹ Jens Volkmann, MD,³⁰ Thorsten Odorfer, MD,³⁰ Andrea A. Kühn, MD,³¹ Friederike Borngreber, MD,³¹ Inke R. König, PhD,³² Carlos Cruchaga, PhD,³³ Adam C. Cotton, BA,³⁴ Gamze Kilic-Berkmen, PhD,³⁴ Alan Freeman, MD,³⁴ Stewart A. Factor, DO,³⁴ Laura Scorr, MD,³⁴ J. Douglas Bremner, MD,^{35,36} Viola Vaccarino, MD, PhD,¹ Arshed A. Quyyumi, MD,³⁷ Christine Klein, MD,²⁴ Joel S. Perlmutter, MD,³⁸ Katja Lohmann, PhD,²⁴ and Hyder A. Jinnah, MD, PhD^{34,39*}

¹Department of Epidemiology, Emory University Rollins School of Public Health, Atlanta, Georgia, USA

²Department of Biomedical Informatics, Emory University School of Medicine, Atlanta, Georgia, USA

³Movement Disorders Division, University of Rochester, Rochester, New York, USA

⁴Neurology One, Orlando, Florida, USA

⁵Department of Neurology, Fixel Institute for Neurological Diseases, University of Florida, Gainesville, Florida, USA

⁶Department of Neurology, University of Maryland School of Medicine, Baltimore, Maryland, USA

⁷Parkinson's Disease Center and Movement Disorders Clinic, Department of Neurology, Baylor College of Medicine, Houston, Texas, USA

⁸Department of Neurology, Henry Ford Health System, Henry Ford Hospital, Detroit, Michigan, USA

⁹Sorbonne Université, Inserm U1127, CNRS UMR 7225, Institut du Cerveau et de la Moelle; Assistance Publique – Hôpitaux de Paris, Hôpital Salpêtrière, Département de Neurologie, Paris, France

¹⁰Department of Neurology, Virginia Commonwealth University, Richmond, Virginia, USA

¹¹Department of Psychology, University of Memphis, Memphis, Tennessee, USA

¹²James J and Joan A Gardner Center for Parkinson's Disease and Movement Disorders, University of Cincinnati Academic Health Center, Cincinnati, Ohio, USA

¹³Booth Gardner Parkinson's Care Center, Evergreen Health, Kirkland, Washington, USA

¹⁴Department of Neurology, University of New Mexico, Albuquerque, New Mexico, USA

¹⁵Beth Israel Deaconess Medical Center, Boston, Massachusetts, USA

¹⁶Department of Neurology, Methodist Neurological Institute, Weill Cornell Medical School, Houston, Texas, USA

¹⁷Icahn School of Medicine at Mount Sinai, Movement Disorders, Department of Neurology, Mount Sinai Beth Israel, New York, New York, USA

¹⁸Unité des troubles du mouvement André-Barbeau, Centre Hospitalier de l'Université de Montréal, Montreal, Canada

¹⁹Department of Neurology, The University of Alabama at Birmingham, Birmingham, Alabama, USA

²⁰Department of Neurology and Psychiatry, Sapienza University of Rome and IRCCS Neuromed, Rome, Italy

²¹Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

²²Neurology, University of California, Davis, Sacramento, California, USA

²³University of Toronto, Edmond J Safra Program in Parkinson Disease; Movement Disorder Clinic, Toronto Western Hospital, Toronto, Ontario, Canada

²⁴Institute of Neurogenetics, University of Lübeck, Lübeck, Germany

²⁵Department of Psychiatry and Psychotherapy, University of Lübeck, Lübeck, Germany

²⁶Department of Neurology, University of Lübeck, Lübeck, Germany

²⁷Institute for Health Sciences, Department of Midwifery Science, University of Lübeck, Lübeck, Germany

*Correspondence to: Dr. Yan V. Sun, Department of Epidemiology, Rollins School of Public Health Emory University 1518 Clifton Rd. NE, Atlanta 30322, GA, USA; E-mail: yan.v.sun@emory.edu and Dr. Hyder A. Jinnah, Department of Neurology, Human Genetics, and Pediatrics, Emory University School of Medicine, 101 Woodruff Circle, Atlanta 30322, GA, USA; E-mail: hjinnah@emory.edu

Relevant conflicts of interest/financial disclosures: None.

Funding agencies: Y.V.S., C.L., Q.H., and H.A.J. are partially supported by the National Institute of Health grant NS096455 and the Dystonia Medical Research Foundation. This work was also supported in part by grants to the Dystonia Coalition, a part of the Rare Diseases Clinical Research Network of the Office of Rare Diseases Research at

the National Center for Advancing Translational Sciences (Grant numbers NS065701, TR001456, and NS116025). K.L., I.R.K., and C.K. are supported by the DFG (FOR2488). The DysTract study has been supported by the Federal Ministry of Education and Sciences (BMBF) in Germany (01GM1514B, to K.L. and C.K.). U.K., M.K., and C.K. were supported by the DFG (TR-SFB134).

Received: 27 August 2019; Revised: 24 June 2021; Accepted: 12 July 2021

Published online in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.28732

²⁸Institute of Systemic Motor Research, University of Lübeck, Lübeck, Germany

²⁹Department of Neurology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

³⁰Department of Neurology, University Hospital Würzburg, Würzburg, Germany

³¹Department of Neurology, Charité Universitätsmedizin Berlin, Berlin, Germany

³²Institute of Medical Biometry and Statistics, University of Lübeck, Lübeck, Germany

³³Department of Psychiatry, Washington University in St. Louis, St. Louis, Missouri, USA

³⁴Department of Neurology, Emory University School of Medicine, Atlanta, Georgia, USA

³⁵Atlanta VA Medical Center, Decatur, Georgia, USA

³⁶Departments of Psychiatry & Behavioral Sciences and Radiology, Emory University School of Medicine, Atlanta, Georgia, USA

³⁷Department of Medicine, Division of Cardiology, Emory University School of Medicine, Atlanta, Georgia, USA

³⁸Department of Neurology, Radiology, Neuroscience, Physical Therapy and Occupational Therapy, Washington University in St. Louis, St. Louis, Missouri, USA

³⁹Department of Human Genetics, Emory University School of Medicine, Atlanta, Georgia, USA

ABSTRACT: Background: Several monogenic causes for isolated dystonia have been identified, but they collectively account for only a small proportion of cases. Two genome-wide association studies have reported a few potential dystonia risk loci; but conclusions have been limited by small sample sizes, partial coverage of genetic variants, or poor reproducibility.

Objective: To identify robust genetic variants and loci in a large multicenter cervical dystonia cohort using a genome-wide approach.

Methods: We performed a genome-wide association study using cervical dystonia samples from the Dystonia Coalition. Logistic and linear regressions, including age, sex, and population structure as covariates, were employed to assess variant- and gene-based genetic associations with disease status and age at onset. We also performed a replication study for an identified genome-wide significant signal.

Results: After quality control, 919 cervical dystonia patients compared with 1491 controls of European ancestry were included in the analyses. We identified one genome-wide

significant variant (rs2219975, chromosome 3, upstream of *COL8A1*, P -value 3.04×10^{-8}). The association was not replicated in a newly genotyped sample of 473 cervical dystonia cases and 481 controls. Gene-based analysis identified *DENND1A* to be significantly associated with cervical dystonia (P -value 1.23×10^{-6}). One low-frequency variant was associated with lower age-at-onset (16.4 ± 2.9 years, P -value = 3.07×10^{-8} , minor allele frequency = 0.01), located within the *GABBR2* gene on chromosome 9 (rs147331823).

Conclusion: The genetic underpinnings of cervical dystonia are complex and likely consist of multiple distinct variants of small effect sizes. Larger sample sizes may be needed to provide sufficient statistical power to address the presumably multi-genic etiology of cervical dystonia. © 2021 International Parkinson and Movement Disorder Society

Key Words: cervical dystonia; genome-wide association study (GWAS); rare disease; movement disorder

The dystonias are a group of disorders characterized by involuntary, repetitive twisting movements, abnormal postures, or other manifestations of excessive muscle contraction.^{1,2} Cervical dystonia (CD) is the most common adult-onset dystonia and is characterized by involuntary patterned contractions of cervical musculature resulting in abnormal movements or postural changes of the head and neck. CD is rare, and estimates of incidence and prevalence are limited by a paucity of comprehensive epidemiological studies.³

Some forms of dystonia are genetically determined, and several genes have been identified so far.⁴⁻⁶ However, these genes collectively account for only a small fraction of all isolated dystonias. Thus, additional genetic and/or environmental factors probably play an important role. The identification of genes among affected pedigrees has been complicated by the fact that many appear to be dominantly inherited

with partial penetrance. Genome-wide association studies (GWAS) agnostically scan markers across the genome in a large population to identify genetic variations associated with a disease trait. GWAS have successfully identified thousands of markers that are associated with a wide spectrum of diseases,⁷ and are the most cost-effective approach to discovering any genetic association with a disease. The power of this method to detect genetic variants depends on study sample size and the underlying genetic architecture. Although variants with large effect sizes can sometimes be detected with sample sizes as small as 100, far more variants with smaller effect sizes can be detected when more than 10,000 cases and controls are examined. Currently, only a few very small GWAS have been published for dystonia. One GWAS involving 212 CD cases from the United Kingdom revealed a genetic variant in *NALCN* gene,⁸ but the

association was not replicated in another study involving additional 252 CD patients from Spain.⁹ A GWAS of 127 cases with musician's dystonia identified a significant association in a locus of *ARSG* gene, but it has not yet been independently replicated.¹⁰

These studies demonstrate that GWAS is a promising approach for the discovery of candidate genes and loci related to the dystonias. However, they are limited by small sample sizes and lack of genome-wide gene-based analysis of low-frequency variants. To address these limitations, the present study provides a comprehensive coverage of the genetic susceptibility in idiopathic CD using a large cohort of 919 cases. Additionally, we investigated potential genetic determinants of age-at-onset and tremor among CD cases.

Patients and Methods

Data Sources and Study Population

The dystonia subjects studied in this project are from the Dystonia Coalition, which is an ongoing study to facilitate collaborations and to advance the pace of clinical and translational research for isolated dystonia syndromes (ie, "primary" dystonias).¹¹⁻¹³ The Dystonia Coalition collected DNA samples and detailed clinical information on adult-onset isolated dystonia patients ($n = 3119$) including several subtypes to identify risk factors and biomarkers to improve prognosis, treatment, and prevention. Because different forms of dystonia may have different genetic substrates, we focused our GWAS on the most common subtype, CD. We included all available CD samples of European ancestry at the time of the study. All cases were evaluated and recruited by individuals with expertise in movement disorders using a standardized protocol, which was videotaped for verification.¹¹ Acquired forms of CD and CD combined with other movement disorders (eg, parkinsonism) were excluded. We included cases who started with dystonia in the neck region (isolated CD). Cases with minor dystonic symptoms in other body regions were also included, if CD was the main problem (segmental or multifocal dystonias).

Control samples without neurological disorders were obtained from the Emory Cardiovascular Biobank study and the Mental Stress-Induced Ischemia, Mechanisms and Prognosis (MIPS). Blood specimens of participants were collected and used to extract genomic DNA. The Dystonia Coalition study and the present GWAS were approved by the Institutional Research Board committee at Emory University. All participants provided written consent. We obtained demographic variables such as age, sex, age-at-onset from the survey and medical records.

Genotypic Measures and Data Processing for the GWAS

Genomic DNA (gDNA) was extracted from blood samples of study participants by a central laboratory at the Coriell Institute for Medical Research (Camden, New Jersey, USA). Each gDNA sample was quantified using the PicoGreen assay, standardized to 50 ng/mL, and processed following the standard Illumina protocol including hybridization, incubation, and scanning. We used Illumina's Multi-Ethnic Genotyping Array (MEGA) platform, which is optimized for genome-wide association studies in multiethnic populations. The MEGA chip directly measures 1.7 million genetic markers, including improved genome-wide coverage of non-European ancestry, exonic content of over 400,000 markers, more than 17,000 variants relevant to clinical and pharmacogenetic studies, and an additional 23,000 variants selected for functional, immunological, oncological, ancestry, forensic, and other common and rare diseases.

Genotypes were measured and called using Illumina's GenomeStudio software. Participants were excluded if they had an overall SNP call rate (ratio of measured SNPs per participant over the total number of SNPs in the data set) $< 95\%$ or mismatch between genotypic and phenotypic measurements (eg, sex and ethnicity). Individuals related closer than second-degree cousins were identified. Only one individual from each related pair was kept in the study. In addition, individual SNPs were excluded from the analyses if they had unknown chromosomal location, were missing for more than 5% of the total participant samples, had Hardy-Weinberg Equilibrium (HWE) P -value less than 10^{-6} among the controls, or had a minor allele frequency (MAF) less than 0.01 among controls.¹⁴ Base-pair position of each variant was annotated to human genome reference GRCh37/hg19.

Cleaned genotype data of common variants were further pruned using plink software with r^2 less than 0.1, and window size and step of 50 and 5 SNPs, respectively. We used ADMIXTURE¹⁵ analysis with the 1000 Genome phase 3 samples and known continental ancestries to assign genotyped individuals to European ($>80\%$ European Ancestry). Principal component analyses were performed using flashPCA software.¹⁶ The dosage data of SNPs in the 1000 Genomes Project phase 3 version 5 reference panel were imputed by the Michigan Imputation Server (<https://imputationserver.sph.umich.edu/>) using minimac.¹⁷ Following imputation, variant-level quality control was performed using the EasyQC R package (www.R-project.org), and exclusion metrics included: Hardy-Weinberg equilibrium P -value $< 10^{-6}$, posterior call probability < 0.9 , imputation quality < 0.3 , MAF < 0.05 , call rate $< 97.5\%$. Variants were also excluded if they deviated $>10\%$ from their expected

allele frequency based on reference data from the Genome Aggregation Database (gnomAD).¹⁸ Among 968 genotyped CD cases with phenotypic data, 13, 14, and 22 samples were excluded due to high missing rate, sex-mismatch, and non-European ancestry, respectively. Following variant-level quality control, we obtained 8,053,795 common variants (MAF > 5%) in 919 CD cases and 1491 controls of European ancestry.

Association Analysis

Genome-wide association analysis of CD status was adjusted for age, sex, and genotype-derived principal components in a logistic regression model. Imputed genetic variants were tested for association with CD assuming an additive genetic model. Among 919 CD patients with available phenotypes, genome-wide association analyses of age-at-onset, tremor, and affected area (segmental vs. focal) were adjusted for age, sex, and genotype-derived principal components in linear or logistic regression models. We summarized the GWAS results using FUMA (<http://fuma.ctglab.nl/>), which is a platform that can be used to annotate, prioritize, visualize, and interpret GWAS results.¹⁹ Genome-wide significant SNPs ($P < 5 \times 10^{-8}$) were grouped into genomic loci if they are dependent on each other at $r^2 > 0.1$ or physically close (distance < 500 kb). Lead SNPs were defined within each locus if they are independent ($r^2 < 0.1$). FUMA uses GWAS summary statistics to compute gene-based P -values using the MAGMA tool.¹⁹ The gene-based P -value is computed for protein-coding genes by mapping SNPs to genes if SNPs are located within the genes. The Bonferroni correction was used to correct for multiple testing based on the total number of tested protein coding genes. The 1000 Genomes phase 3 was used as a reference panel to calculate LD across SNPs and genes. All other statistical analyses were performed in the R statistical environment version 3.2.5 (<http://www.r-project.org/>).

Replication Study

We investigated 473 independent CD patients that originated from the Dystonia Coalition ($n = 270$) or the German Dystonia Registry (DysTract, <http://dystract.cio-marburg.de/en/>, $n = 203$) using a similar protocol for enrollment. We also included 481 healthy controls from Northern Germany (Luebeck area). Variant rs2219975 was genotyped using a melting curve analysis on the LightCycler or by Sanger sequencing.

Results

After quality control, the analysis included 919 cases of CD and 1491 controls, all of European ancestry. The PCA plots showed that the genetic ancestry of the study sample was consistent with the HapMap panel, and the

TABLE 1 Dystonia patients and controls of European ancestry using for the GWAS stage

Variables	Cases (N = 919) Mean ± SD or count (%)	Controls (N = 1491) Mean ± SD or count (%)
Age (y)	60.7 ± 11.4	64.8 ± 11.5
Female	696 (75.7%)	716 (48.0%)
Age at onset (y)	45.6 ± 13.5	N/A
Focal	769 (83.7%)	N/A
Segmental	150 (16.3%)	N/A
Tremor	616 (67.0%)	N/A

Abbreviation: GWAS, genome-wide association study.

genetic background was similar between cases and controls (Fig. S1A,B). Characteristics of these groups are summarized in Table 1. In brief, the CD subjects had a younger average age of 60.7 ± 11.4 years, compared to controls (64.8 ± 11.5 years). More CD participants were females than in the control group (75.7% vs. 48.0%). Among 919 CD subjects, the average age at onset was 45.6 ± 13.5 years.

There was moderate global inflation (inflation factor of 1.07) of over 8 million common variants. Using the 1000 Genome Project-based imputation panel, we identified one genome-wide significant common (MAF > 5%) SNP from a locus on chromosome 3 (Fig. 1 and Table 2). The lead common SNP (rs2219975) of the chromosome 3 locus had a MAF of 0.06 in controls and 0.10 in CD cases. The minor allele C was associated with higher risk of CD (OR of 2.1, P -value = 3.04×10^{-8}). Two variants (chr3:99071470 and chr3:99072830, GRCh37/hg19) with lower MAF (~1%) and high LD in the chromosome 3 locus were also associated with CD with P -values of 6.74×10^{-9} and 8.49×10^{-9} , and ORs of 46 and 50, respectively (see summary statistics of these two low-frequency variants including minor allele counts in Table S1).

The regional plot (Fig. S2) showed all tested SNPs in the neighboring regions of the lead SNP and additional association signals with CD within these regions. The CD-associated locus is located in intergenic region, and is about 300 kb upstream of *COL8A1*.

Genotyping of the lead SNP rs2219975 in the replication samples found the minor C allele less frequently in cases (overall: 6.4%, DC subgroup: 7.4%; DysTract subgroup: 5.2%) compared to controls (9.0%).

Complementary to single variant association test, gene-based statistical methods can analyze multiple genetic markers simultaneously to determine their joint association with a disease trait. Using the MAGMA gene-based analysis method available in the online tool,




FIG. 1. Summary of the genome-wide association study (GWAS) of cervical dystonia. Manhattan plot showed a significant locus on chromosome 3. The horizontal line represented the genome-wide significant threshold of 5×10^{-8} . The quantile-quantile plot showed overall distribution of P -values in the GWAS.

TABLE 2 Summary of common genetic variants ($MAF > 0.05$) of genome-wide significant locus associated with cervical dystonia

SNP ID	rs number	CHR	POS	REF	ALT	AF_Control	AF_CD	OR	T Statistic	P-value
3:99070019	rs2219975	3	99,070,019	T	C	0.06	0.10	2.123	5.539	3.04×10^{-8}

Abbreviations: MAF, minor allele frequency; CHR, chromosome number; POS, base-pair position in GRCh37/hg19; REF, reference allele; ALT, alternative (effective) allele; AF, allele frequency of alternative allele; OR, odds ratio.

functional mapping and annotation (FUMA),¹⁹ the *DENND1A* was significantly associated with CD after correction for multiple testing. We also identified another four genes (Table S2; *ATP11A*, *ST7-OT4*, *PPP1R16B*, *SLC39A12*) with nominal P -values less than 10^{-4} . None of the previously reported genes potentially related to adult-onset isolated dystonia (ie, *ANO3*, *CIZ1*, *COL6A3*, *GNAL*, *THAP1*, *TOR1A*) were significantly associated with CD in the gene-based analysis. We also examined the genetic associations of individual SNPs with CD in regions harboring six previously reported genes related to dystonias, *ANO3*, *CIZ1*, *COL6A3*, *GNAL*, *THAP1*, and *TOR1A*. None of the SNPs in these regions were significantly associated with CD after multiple testing corrections. Gene expression quantitative trait locus (eQTL) analysis showed that eQTLs from the GWAS of CD were the most enriched for brain tissue as well as specific subregions of the brain (eg, cerebellum, cerebellar hemisphere, spinal cord cervical, hippocampus), although not statistically significant after correcting for multiple testing.

There were no common variants significantly associated with age at onset (ranging 8–77 years) among 919 CD

patients adjusted for sex and population structure. Interestingly, one low-frequency SNP on chromosome 9:101222141 (rs147331823, $MAF \sim 0.01$) showed strong association with age at onset (P -value = 3.07×10^{-9}) as shown in Figure S3. The minor allele G was associated with a younger age at onset of 16.4 ± 2.9 years.

The current analyses failed to replicate the CD-associated SNPs in the *NALCN* region on chromosome 13 previously reported.⁸ None of these reported SNPs (reported P -values range from 9.76×10^{-7} to 2.54×10^{-6}) had nominal P -values less than 0.05. We also searched the *NALCN* region including ± 200 kb flanking regions. Among all imputed SNPs in the region, none had a P -value lower than 10^{-3} .

Discussion

The results from this study are derived from the largest GWAS for any type of dystonia to date. CD is considered a rare disorder, and this study was made possible only by pooling samples through a multicenter collaborative effort. However, the sample size of this

study (919 cases and 1491 controls) is considerably smaller than other GWAS for more common disorders such as Parkinson's disease and the finding of a genome-wide significant SNP on chromosome 3 could not be confirmed in a small replication sample. There might be several reasons for the lack of replication: (1) rs2219975 has a rather low frequency (~7%) compared to other genetic risk variants, and thus, larger sample sizes are needed to detect and replicate an association; (2) The frequency of this variant highly depends on genetic ancestry (eg, 0.5481 in Africans and 0.07303 in non-Finnish Europeans, https://gnomad.broadinstitute.org/variant/3-99070019-T-C?dataset=gnomad_r2_1). Small differences in the origin of cases and controls may lead to false positive associations in a GWAS; (3) The individual effect size of genetic risk variants might be low given the known etiological heterogeneity of CD. The sample size was sufficient for genome-wide discovery of CD-related genes with large effects (eg, at alpha level of 5×10^{-8} , OR 1.5 and 1.68 for MAF of 0.2 and 0.1, respectively), but insufficient for genetic associations with small effects or very low-risk allele frequency.

In the present GWAS of CD, we could not confirm a previously reported association of CD in *NALCN* from an independent GWAS. We also were unable to replicate the association of dystonia with the *ARSG* gene, although the latter may reflect the different type of dystonia studied (eg, task specific dystonia).¹⁰

The locus on chromosome 3 is located upstream of *COL8A1*. The *COL8A1* gene encodes one of the two alpha chains of type VIII collagen. Defects in *COL8A1* are associated with corneal dystrophy and age-related macular degeneration. It has not been associated with any type of dystonia, but it is interesting to note that dystonia has been reported to be potentially caused by biallelic mutations in *COL6A3*, another member of the collagen family.²⁰

Gene-based analysis identified *DENND1A* on chromosome 9 to be significantly associated with CD. *DENND1A* encodes DENN domain containing protein 1A, a member of the connectenn family. It serves as a guanine nucleotide exchange factor that is expressed in brain and plays a role in vesicle function. It has not yet been associated with dystonia, although pathogenic variants in different guanine nucleotide exchange factors encoded by *GNAL* and *GNAO1* have been linked to adult onset focal and segmental dystonias and a childhood-onset movement disorder that combines chorea and dystonia,⁴ respectively. *SLC39A12* (solute carrier family 39 member 12) encodes a zinc transporter and belongs to a subfamily of proteins that share structural characteristics of zinc transporters. Mutations in *SLC39A14*, another member of the same ZIP transporter family, have been linked to hypermanganesemia associated with infantile-onset dystonia and parkinsonism-dystonia.^{21,22}

Although we did not identify any genome-wide significant common variant associated with age at onset for

CD, a low-frequency variant (MAF ~1%) was associated with a decrease in age at onset (16.4 ± 2.9 years, P -value = 3.07×10^{-8}), located within the *GABBR2* gene on chromosome 9. This gene encodes a subunit of GABA-B receptors, which play an important role in neuronal excitability. This gene has not been previously associated with dystonia, although GABA receptor activators such as baclofen and benzodiazepines can be effective in treating some types of dystonia. Because the strong association was based on nine carriers of the minor allele, future association study is warranted to validate the finding.

In conclusion, we have conducted the largest GWAS of any type of dystonia so far assembled, and attempted to address heterogeneity by relying on a standardized exam and by focusing on a single dystonia subtype, CD. Despite these measures, GWAS did not reveal any robust locus, indicating a presumably rather small impact of an individual common variant at the population level. Future genetic studies of dystonias require the whole scientific community to join effort to reveal the complex genetic architecture of CD and other dystonias. ■

Data Availability Statement

Data sharing is not applicable to this article as no new data were created or analyzed in this study. GWAS data of cervical dystonia are shared via dbGaP Study Accession: phs001803.v1.p1.

References

- Albanese A et al. Phenomenology and classification of dystonia: a consensus update. *Mov Disord* 2013;28:863–873.
- Jinnah HA. The Dystonias. *Continuum (Minneapolis)* 2019; 25(4):976–1000.
- Defazio G, Berardelli A, Hallett M. Do primary adult-onset focal dystonias share aetiological factors? *Brain* 2007;130(Pt 5):1183–1193.
- Jinnah HA, Sun YV. Dystonia genes and their biological pathways. *Neurobiol Dis* 2019;129:159–168.
- Balint B, Bhatia KP. Isolated and combined dystonia syndromes - an update on new genes and their phenotypes. *Eur J Neurol* 2015; 22(4):610–617.
- Balint B et al. Dystonia. *Nat Rev Dis Primers* 2018;4(1):25.
- Welter D et al. The NHGRI GWAS catalog, a curated resource of SNP-trait associations. *Nucleic Acids Res* 2014;42(Database issue): D1001–D1006.
- Mok KY et al. Genomewide association study in cervical dystonia demonstrates possible association with sodium leak channel. *Mov Disord* 2014;29(2):245–251.
- Gomez-Garre P et al. Lack of validation of variants associated with cervical dystonia risk: a GWAS replication study. *Mov Disord* 2014; 29(14):1825–1828.
- Lohmann K et al. Genome-wide association study in musician's dystonia: a risk variant at the arylsulfatase G locus? *Mov Disord* 2014; 29(7):921–927.
- Yan L et al. Secured web-based video repository for multicenter studies. *Parkinsonism Relat Disord* 2015;21(4):366–371.
- LeDoux MS et al. Clinical and genetic features of cervical dystonia in a large multicenter cohort. *Neurol Genet* 2016;2(3):e69.
- Kilic-Berkmen G et al. The dystonia coalition: a multicenter network for clinical and translational studies. *Front Neurol* 2021;12:415.

14. Anderson CA et al. Data quality control in genetic case-control association studies. *Nat Protoc* 2010;5(9):1564–1573.
15. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated individuals. *Genome Res* 2009;19(9):1655–1664.
16. Tsai TH et al. LC-MS/MS-based serum proteomics for identification of candidate biomarkers for hepatocellular carcinoma. *Proteomics* 2015;15(13):2369–2381.
17. Das S et al. Next-generation genotype imputation service and methods. *Nat Genet* 2016;48(10):1284–1287.
18. Karczewski KJ et al. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature* 2020;581(7809):434–443.
19. Watanabe K et al. Functional mapping and annotation of genetic associations with FUMA. *Nat Commun* 2017;8(1):1826.
20. Lohmann K et al. The role of mutations in COL6A3 in isolated dystonia. *J Neurol* 2016;263(4):730–734.
21. Tuschl K et al. Mutations in SLC39A14 disrupt manganese homeostasis and cause childhood-onset parkinsonism-dystonia. *Nat Commun* 2016;7:11601.
22. Juneja M et al. A novel mutation in SLC39A14 causing hypermanganesemia associated with infantile onset dystonia. *J Gene Med* 2018;20(4):e3012.

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.

SGML and CITI Use Only DO NOT PRINT

Author Roles

Y.V.S. and H.A.J. were involved in the conception, organization, and execution of the research project; R.B., R.R., I.M., S.R., K.B., K.H., J.J., N.J.P., E.R., M.V., B.B., M.S.L., A.E., P.A., S.P., S.A.F., W.G.O., R.S., S.C., S.N., A.B., A.P., A.B., S.H.F., M.K., U.K., M.N., T.B., S.L., M.B., S.Z., A.M., M.G., J.V., T.O., A.A.K., F.B., A.C., G.K., A.F., S.F., L.S., J.D.B., V.V., A.A.Q., C.K., J.P., K.L., and H.A.J. contributed to evaluation of cases, and data collection and biospecimen collection; Y.V.S., C.L., I.R.K., and Q.H. were responsible for the design and execution of statistical analysis; The first draft was written by Y.V.S. and H.A.J., and developed further by C.K., K.L., M.S.L., and W.G.O. All coauthors contributed to critical review and editing of the manuscript.

Full Financial Disclosures for the past 12 Months

P.A. has received honoraria from Sunovion, Acadia, Amneal, Accordia, ADAMA, Kyowa Kirin, and Supernus for speaking. She has also received consulting fees from ADAMAS CALA Health, and Sunovion. Research support has been received from US WorldMeds, Centogene, Biogen, Merck, Sun Pharma, Advanced Research Company Limited (SPARC) Anexxon Inc, CHDI Foundation, Roche.

T.B. has received grant support from DFG FOR 2698. He has also received honoraria or consulting fees from Ipsen Pharma, Allergan, Merz Pharmaceuticals.

B.D.B. has received research grant support from the Parkinson's Foundation, Dana Foundation, NIH (NIH/NCATS Colorado CTSI Grant Number KL2 TR001080), Dystonia Coalition (receives the majority of its support through NIH grant NS065701 from the Office of Rare Diseases Research in the National Center for Advancing Translational Science and National Institute of Neurological Disorders and Stroke), and from Mary Rossick Kern and Jerome H. Kern. He also serves on the medical advisory boards for the Benign Essential Blepharospasm Research Foundation and the National Spasmodic Torticollis Association.

F.B. has received grant support from Clinician Scientist Program, BIH, Charité-Universitätsmedizin Berlin, and Max-Delbrück Centre.

M.B. has received honoraria from Ipsen Pharma.

A.B. has active or recent grant support from the NIH Revance, Allergan, Merz, Ipsen. She has served as a consultant or on advisory boards for Revance, Ipsen, Allergan, and Merz. She is a board member for the American Board of Psychiatry and Neurology, California Institute of Regenerative Medicine, McKnight Brain Research Foundation, Latinx Physicians of California. She is founder and board member of Care Directions.

C.C. has received research support from Biogen, Eisai, and Alector. He has consulted or is a member of the advisory board of Vivid genetics, Halia Therapeutics and Takeda and GSK.

A.J.E. has received grant support from the NIH and the Michael J Fox Foundation; personal compensation as a consultant/scientific advisory board member for Abbvie, Neuroderm, Neurocrine, Amneal, Acadia, Acorda, Kyowa Kirin, Sunovion, Lundbeck, and USWorldMeds; honoraria from Acadia, Sunovion, USWorldMeds; and publishing royalties from Lippincott Williams & Wilkins, Cambridge University Press, and Springer.

S.A.F. has honoraria from Lundbeck, Teva, Sunovion, Biogen, Acadia, Neuroderm, Acorda, CereSpir; grants from Ipsen, Medtronic, Boston Scientific, Teva, US World Meds, Sunovion Therapeutics, Vaccinex, Voyager, Jazz Pharmaceuticals, Lilly, CHDI Foundation, Michael J. Fox Foundation, NIH; Royalties from Demos, Blackwell Futura, Springer for textbooks, Uptodate; and other support from Bracket Global LLC, CNS Ratings LLC.

S.H.F. has received Clinic support from the Edmond J Safra Foundation for Parkinson Research; National Parkinson Foundation and the Toronto Western and General Foundation. Salary from UHN Dept of Medicine Practice Plan. Research funding from Michael J Fox Foundation for Parkinson Research, NIH (Dystonia Coalition); CIHR; Parkinson Canada. Honoraria from the International Parkinson and Movement Disorder Society and American Academy of Neurology. Site PI for Clinical Trials for Biotie, Cynapsus, Eisai, Revance. Consultancy/speaker fees from Acadia, Atuka, Sunovion, Teva, Paladin; royalties from Oxford University Press.

M.G. has received grant support from Merz Pharmaceuticals, Allergan, DFG, Werner Otto Stiftung, Ostseeförderung. He has served on advisory boards for Merz Pharmaceuticals and Allergan, and has received honoraria from Merz Pharmaceuticals, Allergan, and Ipsen.

H.A.J. has active or recent grant support from the US government (National Institutes of Health), private philanthropic organizations (Cure Dystonia Now), and industry (Cavion Therapeutics, Retrophin Inc., Revance Therapeutics). He has also served on advisory boards or as a consultant for Abide Therapeutics, Allergan Inc., CoA Therapeutics, Retrophin Inc, and Revance Therapeutics. He has received honoraria or stipends for lectures or administrative work from the International Parkinson's Disease and Movement Disorders Society. He serves on the scientific advisory boards for several private foundations including the Benign Essential Blepharospasm Research Foundation, Cure Dystonia Now, the Dystonia Medical Research Foundation, the Tourette Association of American, and Tyler's Hope for a Cure. He also is director of the Dystonia Coalition, which receives the majority of its support through NIH grants (NS116015, NS067501, and TR001456 from the National Institutes of Neurological Disorders and Stroke and the Office of Rare Diseases Research at the National Center for Advancing Translational Sciences). The Dystonia Coalition has received additional material or administrative support from industry sponsors (Allergan Inc. and Merz Pharmaceuticals) as well as private foundations (the American Dystonia Society, Beat Dystonia, the Benign Essential Blepharospasm Foundation, Cure Dystonia Now, Dystonia Europe, Dystonia Inc., Dystonia Ireland, the Dystonia Medical Research Foundation, the Foundation for Dystonia Research, the National Spasmodic Dysphonia Association, and the National Spasmodic Torticollis Association).

M.K. has received grant support from TCRC-134, FOR 2488.

C.K. has received grant support from the Hermann and Lilly Schilling Foundation; the German Research Foundation; the BMBF; the German Research Foundation; the European Community; intramural funds from the University of Luebeck. She has served as medical advisor to Centogene. She has also received honoraria from the Else Kroener Fresenius Foundation.

I.R.K. has received grant support from the German Research Foundation, BMBF, German Cancer Aid.

U.M.K. has received grant support from the German Research Foundation.

A.A.K. has received honoraria or served on advisory boards for Medtronic und Boston Scientific.

M.S.L. serves on the speaker's bureaus for Acadia, Teva, Acorda and Adamas, and editorial boards of *Neurology* and *Tremor and Other Hyperkinetic Movements*; has received research support from the National Institutes of Health, Dystonia Medical Research Foundation, Department of Defense, Michael J. Fox Foundation, Revance, and PharmaTwoB; and receives royalty payments from Elsevier for editing *Animal Models of Movement Disorders* and *Movement Disorders: Genetics and Models*.

S.L. has nothing to disclose.

K.L. has received grant support from German Research Foundation; Federal Ministry of Education and Science (BMBF); Movement Disorder Society; Damp-Stiftung.

A.M. has nothing to disclose.

W.G.O., MD has received honorarium for speaking bureau from Teva, ACADIA, Acorda, Neurocrine, UCBPharma, USWorldMeds, and Lundbeck. He has received research grants from Biogen, Lundbeck, Sun, Restless Legs Syndrome Foundation, Parkinson's Study Group, Lilly, and Revance. He receives royalties from the books *Restless Legs Syndrome*, *Movement Disorders in Clinical Practice*, and *UpToDate*.

M.N. has nothing to disclose.

T.O. has received honoraria or travel grants from Allergan, Teva, and Merz.

E.R. received research support from Merz-Pharma, Orkyn, Aguettant, Elivie, Ipsen, Fondation Desmarest, AMADYS, Agence Nationale de la Recherche, and Fonds de Dotation Brou de Laurière; has served on scientific advisory boards for Orkyn, Aguettant, Merz-Pharma and has received speech honoraria from Orkyn, Aguettant, Merz-Pharma, Medday-Pharma, Everpharma, International Parkinson and Movement disorders Society; received travel grant from Vitalair, Aguettant, Merz-Pharma, Ipsen, Merck, Orkyn, Elivie, Dystonia medical Research Foundation, International Parkinson and Movement disorders Society, European Academy of Neurology, International Association of Parkinsonism and Related Disorders.

J.V. has received grant support from Boston Scientific, Medtronic, BMBF, Deutsche Forschungsgemeinschaft. He has received honoraria from Boston Scientific, Medtronic, BMBF, Deutsche Forschungsgemeinschaft. He has served as a consultant or on an advisory board for Boston Scientific, Newronika, and Medtronic.

S.Z. has received grant support from the European Union and Werner Otto Foundation and honoraria from Merz Pharmaceuticals and Brainlab AG.