

Henry Ford Health

Henry Ford Health Scholarly Commons

Surgery Articles

Surgery

3-1-2023

ASO Visual Abstract: Multiple Pretransplant Treatments for Patients Without Pathological Complete Response may Worsen Posttransplant Outcomes in Patients With Hepatocellular Carcinoma

Shingo Shimada

Henry Ford Health, sshimad1@hfhs.org

Tayseer Shamaa

Henry Ford Health, tshamaa1@hfhs.org

Tommy Ivanics

Henry Ford Health, tivanic1@hfhs.org

Toshihiro Kitajima

Henry Ford Health, tkitaji1@hfhs.org

Mohamed Adhnan

Henry Ford Health

See next page for additional authors

Follow this and additional works at: https://scholarlycommons.henryford.com/surgery_articles

Recommended Citation

Shimada S, Shamaa T, Ivanics T, Kitajima T, Adhnan M, Collins K, Rizzari M, Yoshida A, Abouljoud M, Salgia R, and Nagai S. ASO Visual Abstract: Multiple Pretransplant Treatments for Patients Without Pathological Complete Response may Worsen Posttransplant Outcomes in Patients With Hepatocellular Carcinoma. *Ann Surg Oncol* 2022.







This Article is brought to you for free and open access by the Surgery at Henry Ford Health Scholarly Commons. It has been accepted for inclusion in Surgery Articles by an authorized administrator of Henry Ford Health Scholarly Commons.

Authors

Shingo Shimada, Tayseer Shamaa, Tommy Ivanics, Toshihiro Kitajima, Mohamed Adhnan, Kelly Collins, Michael Rizzari, Atsushi Yoshida, Marwan S. Abouljoud, Reena Salgia, and Shunji Nagai




ASO Visual Abstract: Multiple Pretransplant Treatments for Patients Without Pathological Complete Response may Worsen Posttransplant Outcomes in Patients With Hepatocellular Carcinoma

Shingo Shimada, MD, PhD¹ , Tayseer Shamaa, MD¹, Tommy Ivanics, MD, MPH¹ ,
Toshihiro Kitajima, MD, PhD¹ , Mohamed Adhnan, MD¹, Kelly Collins, MD¹ ,
Michael Rizzari, MD¹, Atsushi Yoshida, MD¹, Marwan Abouljoud, MD¹ , Reena Salgia, MD², and
Shunji Nagai, MD, PhD¹ 

¹Division of Transplant and Hepatobiliary Surgery, Henry Ford Health System, Detroit, MI; ²Division of Gastroenterology and Hepatology, Henry Ford Health System, Detroit, MI

In this study (<https://doi.org/10.1245/s10434-022-12789-2>), we revealed a history of multiple pre-LT treatments without pCR was associated with risk of post-LT recur-

rence of HCC. Also, Y90 was possible treatment most likely to be associated with achieving pCR.



FUNDING Nothing to disclose. This work received no financial support.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

DISCLOSURE The authors of this manuscript have no conflicts of interest to disclose.