

Henry Ford Health

## Henry Ford Health Scholarly Commons

---

Cardiology Articles

Cardiology/Cardiovascular Research

---

2-1-2021

### Determinants of Hospital Variation in Cardiac Rehabilitation Enrollment During Coronary Artery Disease Episodes of Care

Michael P. Thompson

Jessica M. Yaser

Hechuan Hou

John D. Syrjamaki

Alphonse DeLucia

*See next page for additional authors*

Follow this and additional works at: [https://scholarlycommons.henryford.com/cardiology\\_articles](https://scholarlycommons.henryford.com/cardiology_articles)

---

#### Recommended Citation

Thompson MP, Yaser JM, Hou H, Syrjamaki JD, DeLucia A, 3rd, Likosky DS, Keteyian SJ, Prager RL, Gurm HS, and Sukul D. Determinants of Hospital Variation in Cardiac Rehabilitation Enrollment During Coronary Artery Disease Episodes of Care. *Circ Cardiovasc Qual Outcomes* 2021; 14(2):e007144.

This Article is brought to you for free and open access by the Cardiology/Cardiovascular Research at Henry Ford Health Scholarly Commons. It has been accepted for inclusion in Cardiology Articles by an authorized administrator of Henry Ford Health Scholarly Commons.

---

## Authors

Michael P. Thompson, Jessica M. Yaser, Hechuan Hou, John D. Syrjamaki, Alphonse DeLucia, Donald S. Likosky, Steven J. Keteyian, Richard L. Prager, Hitinder S. Gurm, and Devraj Sukul

ORIGINAL ARTICLE

# Determinants of Hospital Variation in Cardiac Rehabilitation Enrollment During Coronary Artery Disease Episodes of Care

**BACKGROUND:** Cardiac rehabilitation (CR) is associated with improved outcomes for patients with coronary artery disease (CAD). However, CR enrollment remains low and there is a dearth of real-world data on hospital-level variation in CR enrollment. We sought to explore determinants of hospital variability in CR enrollment during CAD episodes of care: medical management of acute myocardial infarction (AMI-MM), percutaneous coronary intervention (PCI), and coronary artery bypass grafting (CABG).

**METHODS:** A cohort of 71 703 CAD episodes of care were identified from 33 hospitals in the Michigan Value Collaborative statewide multipayer registry (2015 to 2018). CR enrollment was defined using professional and facility claims and compared across treatment strategies: AMI-MM (n=18 678), PCI (n=41 986), and CABG (n=11 039). Hierarchical logistic regression was used to estimate effects of predictors and hospital risk-adjusted rates of CR enrollment.

**RESULTS:** Overall, 20 613 (28.8%) patients enrolled in CR, with significant differences by treatment strategy: AMI-MM=13.4%, PCI=29.0%, CABG=53.8% ( $P<0.001$ ). There were significant differences in CR enrollment across age groups, comorbidity status, and payer status. At the hospital-level, there was over 5-fold variation in hospital risk-adjusted CR enrollment rates (9.8%–51.6%). Hospital-level CR enrollment rates were highly correlated across treatment strategy, with the strongest correlation between AMI-MM versus PCI ( $R^2=0.72$ ), followed by PCI versus CABG ( $R^2=0.51$ ) and AMI-MM versus CABG ( $R^2=0.46$ , all  $P<0.001$ ).

**CONCLUSIONS:** Substantial variation exists in CR enrollment during CAD episodes of care across hospitals. However, within-hospital CR enrollment rates were significantly correlated across all treatment strategies. These findings suggest that CR enrollment during CAD episodes of care is the product of hospital-specific rather than treatment-specific practice patterns.

Michael P. Thompson<sup>1</sup>, PhD

Jessica M. Yaser, MPH

Hechuan Hou<sup>2</sup>, MS

John D. Syrjamaki, MS

Alphonse DeLucia III, MD

Donald S. Likosky<sup>3</sup>, PhD

Steven J. Keteyian, PhD

Richard L. Prager, MD

Hitinder S. Gurm, MBBS

Devraj Sukul, MD, MS

**Key Words:** cardiac rehabilitation  
■ coronary artery disease ■ episode of care ■ hospital ■ quality of care

© 2021 American Heart Association, Inc.

<https://www.ahajournals.org/journal/circoutcomes>

## WHAT IS KNOWN

- Cardiac rehabilitation (CR) confers significant benefit to patients hospitalized with coronary artery disease, including those receiving medical management for acute myocardial infarction, percutaneous coronary intervention, or coronary artery bypass grafting.
- Enrollment in CR remains low and variable across hospitals, but the underlying determinants of enrollment are not well known.

## WHAT THE STUDY ADDS

- CR enrollment rates after coronary artery disease admissions varied by 5-fold across hospitals but within-hospital enrollment rates across treatment strategies were similar.
- The effect of payer and admitting hospital on CR enrollment was as large as or greater than patient demographic and clinical factors.
- Patients in high CR enrollment hospitals attended more sessions within 90 days of discharge with fewer days from discharge to first CR session.

Cardiac rehabilitation (CR) may confer significant clinical benefit for patients with coronary artery disease (CAD), including lower cardiovascular mortality, fewer readmissions, and improved quality of life.<sup>1,2</sup> Guidelines strongly recommend CR for patients who have undergone coronary artery bypass grafting (CABG) or percutaneous coronary intervention (PCI), as well as patients medically managed for acute myocardial infarction (AMI-MM).<sup>3–6</sup> While there has been considerable focus on increasing CR referral rates, wide geographic variability persists, and only 20% to 40% of eligible patients ultimately enroll in CR.<sup>7,8</sup>

Despite the demonstrated benefits of CR, few eligible patients receive a CR referral and even fewer patients ultimately enroll in a single CR session.<sup>9–12</sup> Improving CR enrollment during episodes of care for CAD at a hospital-level requires a better understanding of the variation and underlying determinants of enrollment. Prior studies have focused primarily on referral to CR and have shown wide variability in referral rates across hospitals.<sup>9</sup> However, referral rates are an inadequate proxy for enrollment, with reported referral rates >2-fold higher than enrollment.<sup>8</sup> Moreover, the extent to which variation in CR enrollment is a product of patient complexity or hospital-specific practice patterns is not known. Even less is known about CR enrollment across CAD treatment strategies, as clinical registries are often procedure-based or do not capture post-discharge CR enrollment.<sup>13</sup> Identifying determinants of CR enrollment variation may help to inform hospital-level quality improvement initiatives seeking to boost enrollment after CAD events.

In this context, we sought to explore hospital-level variation in CR enrollment during CAD episodes of care by leveraging a statewide, multipayer administrative claims registry in Michigan. Specifically, we sought to identify determinants of variation in CR enrollment across hospitals, the extent to which variation in risk-adjusted CR enrollment can be attributed to the hospital level, and within-hospital variation in CR enrollment by treatment strategy: AMI-MM, PCI, and CABG. We hypothesize that CR enrollment will vary widely across hospitals, with substantial variation in enrollment attributed to the hospital-level.

## METHODS

This study was deemed exempt from human subjects' protections by the University of Michigan Institutional Review Board. Members of the study team (Dr Thompson, J.M. Yaser, J.D. Syrjamaki) had direct access to the data throughout the study and take responsibility for its integrity and the data analysis. Because of the sensitive nature of the data in this study, public access to the data will not be provided.

## Data Source and Sample

Data from the Michigan Value Collaborative (MVC) administrative claims data registry were used for this study. The MVC is a statewide collaborative quality initiative representing 87 Michigan hospitals and 39 provider organizations supported by Blue Cross Blue Shield of Michigan (BCBSM) and Blue Care Network (BCN) with the goal of helping Michigan hospitals achieve the best possible outcomes at the lowest reasonable cost. The MVC registry contains facility and professional administrative claims data for all Michigan residents insured by BCBSM/BCN commercial plans, BCBSM/BCN Medicare Advantage plans and Medicare fee-for-service (FFS). Administrative claims are arranged into episodes of care, including all facility and professional claims provided during the index admission or procedure and within 90 days of discharge, which have been validated against external data sources.<sup>14</sup> Details on the construction and definitions of MVC registry episodes of care can be found in the Appendix.

The MVC registry contains episodes of care for AMI-MM, PCI, and CABG, which were defined using *International Classification of Diseases (ICD)* versions 9 and 10 diagnosis and procedure codes from facility claims and Current Procedure Terminology (CPT) codes from professional claims (Table 1 in the [Data Supplement](#)). We note that PCI episodes of care include both inpatient and outpatient procedures. Episodes from the MVC registry were eligible for inclusion in the study if they (1) had an index admission or procedure date between January 1, 2015 and December 31, 2018 and (2) were treated at one of the 33 Michigan hospitals that provide both PCI and CABG services. Patients were excluded if they died during the index admission or procedure, were discharged to hospice care, or had missing discharge disposition information.

## CR Enrollment

Current procedural terminology (CPT) codes and health care common procedure codes were used to identify claims for CR in facility and professional claims (CPT: 93797 and 93798,

health care common procedure codes: G0422 and G0423, revenue center code: 943).<sup>15</sup> A binary variable was created indicating any CR enrollment within 90 days of discharge (yes versus no). Secondary CR enrollment variables included the number of days from discharge to the first CR visit and the number of CR sessions attended within 90 days of discharge.

## Covariates


Several patient covariates were included in the analysis, including demographics, clinical factors, and

comorbidities. Demographics included patient age category (85+ years, 75–84 years, 65–74 years, 55–64 years, and <55 years), sex (male versus female), and payer type (BCBSM/BCN Commercial, BCBSM/BCN Medicare Advantage and Medicare FFS). Clinical factors included 6-month prior spending above the median for the sample, treatment strategy (AMI-MM, PCI, or CABG), and the presence of comorbidities within 6 months of admission. Comorbidities occurring in the 6 months before admissions were derived from administrative claims data

**Table 1. Patient Characteristics by CR Enrollment Status**

Patient characteristics	Overall	CR enrollment		P value
		No	Yes	
Sample, n	71 703 (100)	51 090 (71)	20 613 (29)	...
Hospitals, N	33	33	33	...
Male, n (%)	45 696 (63.7)	31 352 (61.4)	14 344 (69.6)	<0.001
Age category, n (%)				
85+ y	7989 (11.1)	7126 (14.0)	863 (4.2)	<0.001
75–84 y	18 710 (26.1)	13 613 (26.7)	5097 (24.7)	...
65–74 y	26 092 (36.4)	17 525 (34.3)	8567 (41.6)	...
55–64 y	12 929 (18.0)	8741 (17.1)	4188 (20.3)	...
<55 y	5983 (8.3)	4085 (8.0)	1898 (9.2)	...
Large 6-mo prior spending, n (%)	35 371 (49.3)	26 288 (51.5)	9083 (44.1)	<0.001
Payer, n (%)				
Medicare FFS	42 547 (59.3)	31 694 (62.0)	10 853 (52.7)	<0.001
Medicare advantage	8336 (11.6)	6026 (11.8)	2310 (11.2)	...
Commercial	20 820 (29.0)	13 370 (26.2)	7450 (36.1)	...
Admission type, n (%)				
AMI-MM	18 678 (26.0)	16 167 (31.6)	2511 (12.2)	<0.001
PCI	41 986 (58.6)	29 817 (58.4)	12 169 (59.0)	...
CABG	11 039 (15.4)	5106 (10.0)	5933 (28.8)	...
Number of HCCs, mean (SD)				
Mean (SD)	5(3)	5 (3)	4 (2)	<0.001
Median (IQR)	4 (4)	4 (4)	3 (3)	
Major HCC categories, n (%)				
Diabetes	33 426 (46.6)	24 795 (48.5)	8631 (41.9)	<0.001
Congestive heart failure	27 217 (38.0)	21 812 (42.7)	5405 (26.2)	<0.001
Vascular disease	22 316 (31.1)	17 440 (34.1)	4876 (23.7)	<0.001
COPD	18 699 (26.1)	14 987 (29.3)	3712 (18.0)	<0.001
Cancer	7869 (11.0)	5747 (11.3)	2122 (10.3)	0.0002
Neurological disorders	5340 (7.5)	4313 (8.4)	1027 (5.0)	<0.001
Chronic kidney disease	4847 (6.8)	4261 (8.3)	586 (2.8)	<0.001
Psychiatric disorder	4542 (6.3)	3583 (7.0)	959 (4.7)	<0.001
Respiratory dysfunction	3297 (4.6)	2611 (5.1)	686 (3.3)	<0.001
Cerebrovascular disease	3134 (4.4)	2551 (5.0)	583 (2.8)	<0.001
Drug/alcohol disorders	1415 (2.0)	1135 (2.2)	280 (1.4)	<0.001
Chronic liver disease	1148 (1.6)	941 (1.8)	207 (1.0)	<0.001

AMI-MM indicates acute myocardial infarction medically managed; CABG, coronary artery bypass grafting; COPD, chronic obstructive pulmonary disease; CR, cardiac rehabilitation; FFS, fee-for-service; HCC, hierarchical condition category; IQR, interquartile range; and PCI, percutaneous coronary intervention.




**Figure 1.** Quarterly trends in cardiac rehabilitation enrollment from 2015 to 2018 for medically managed acute myocardial infarction (AMI-MM), percutaneous coronary intervention (PCI), and coronary artery bypass grafting (CABG). CR indicates cardiac rehabilitation.

using established methods to develop 79 hierarchical condition categories (HCCs), which were grouped into 12 major HCC categories for analysis (Table II in the [Data Supplement](#)).<sup>16,17</sup> The mean and SD number of comorbidities was estimated for each patient. Additionally, the presence of AMI (yes or no) was included as a covariate in the PCI and CABG episodes of care. Each patient was also attributed to its admitting hospital according to its national provider identifier.

Statistical Analysis

Significant differences in patient covariates across CR enrollment status were evaluated in bivariate analyses. Chi-square tests and ANOVA were used to test for differences in categorical and continuous patient covariates, respectively. Quarterly trends in CR enrollment were plotted over time, stratified by treatment strategy (AMI-MM, PCI, and CABG). Rates of CR enrollment were also compared across payers for all CAD events and by treatment strategy using  $\chi^2$  tests.



**Figure 2.** Ninety-day cardiac rehabilitation (CR) enrollment by payer and treatment strategy: medically managed acute myocardial infarction (AMI-MM), percutaneous coronary intervention (PCI), and coronary artery bypass grafting (CABG). CAD indicates coronary artery disease; and FFS, fee-for-service.

Next, hierarchical logistic regression modeling was used to estimate the independent association between patient covariates and CR enrollment, and the model variance for hospital random effects. Four separate models of CR enrollment were created, including a model for all CAD episodes of care, and stratified by treatment strategy: AMI-MM, PCI, and CABG. All models were adjusted for sex, age category, large 6-month prior spending, payer, and major HCC categories. Additionally, the primary model for all CAD episodes of care adjusted for treatment strategy. In the models for PCI and CABG, additional adjustment included whether or not the patient was admitted with AMI (yes versus no).

From these models, the expected probability each patient received CR was estimated and then summed at the hospital level to sum the expected number of patients who received CR. The ratio of observed to expected (ie, the O/E ratio) number of patients receiving CR for each hospital was multiplied by the overall CR enrollment rate to calculate the hospital-level risk-adjusted CR enrollment rate. A dot plot was used to illustrate the variation in CR enrollment across hospitals for all CAD episodes of care, and by treatment strategy (AMI-MM, PCI, CABG). The hospital random effect variance was estimated from each hierarchical regression model and used to estimate the intraclass correlation coefficient (ICC) for the hospital. The ICC estimates the percent of the variation in CR enrollment that can be attributed to the hospital-level, adjusted for patient factors.<sup>18</sup> The hospital random effect variance was also used to estimate the median odds ratio (MOR), which is a measure of variation that expresses the relative odds of CR enrollment for 2 identical patients discharged from one randomly selected hospital to a second randomly selected hospital. As a sensitivity analysis, the analysis was repeated in only patients who were discharged to home (ie, excluding patients who were discharged to skilled nursing facilities, inpatient rehabilitation, or other extended care facilities).

To assess within-hospital CR enrollment across treatment strategy, scatter plots compared risk-adjusted CR enrollment with pairwise comparisons for AMI-MM, PCI, and CABG. The  $R^2$  for each pairwise comparison was estimated to assess the extent to which variation in CR enrollment is explained across treatment strategies.

Finally, hospitals were placed into tertiles according to their rank-order in risk-adjusted CR enrollment for all CAD episodes of care, and by treatment strategy, representing low, moderate, and high CR enrollment. We compared the frequency and percent of any CR enrollment across tertiles, the mean (SD) number of sessions, and mean (SD) days to the first session among those who were enrolled in CR. Trends in these variables across hospital CR enrollment tertiles were tested using the Cochran-Armitage test for trend and for trends in mean number of sessions and days to first session across tertiles using ANOVA.

All analyses were performed using SAS 9.4 and Stata 15.1, and statistical tests were deemed significant at an alpha of 0.05 (2-sided).

## RESULTS

A total of 71 703 discrete CAD episodes of care were identified between January 2015 and December 2018,


**Table 2. Multivariable Model of Patient Characteristics Associated With CR Enrollment During CAD Episodes of Care**

Patient characteristic	Odds ratio (95% CI)	P value
Female vs male	0.93 (0.90–0.97)	0.0004
Age		
85+	0.50 (0.45–0.55)	<0.001
75–84	1.15 (1.07–1.24)	0.0002
65–74	1.35 (1.26–1.45)	<0.001
55–64	1.08 (1.01–1.16)	0.027
<55	1 (Ref)	...
Large 6-mo prior spending	1.06 (1.02–1.10)	0.004
Payer		
Medicare FFS	0.70 (0.67–0.73)	<0.001
Medicare advantage	0.70 (0.66–0.75)	<0.001
Commercial	1 (Ref)	...
Treatment strategy, n (%)		
AMI-MM	0.47 (0.45–0.49)	<0.001
PCI	1 (Ref)	...
CABG	2.74 (2.62–2.87)	<0.001
Major HCC categories		
Cancer	1.05 (0.99–1.11)	0.130
Cerebrovascular disease	0.81 (0.73–0.90)	<0.001
Chronic kidney disease	0.50 (0.46–0.56)	<0.001
Chronic liver disease	0.70 (0.59–0.82)	<0.001
COPD	0.68 (0.65–0.71)	<0.001
Congestive heart failure	0.72 (0.69–0.75)	<0.001
Diabetes	0.83 (0.80–0.87)	<0.001
Drug/alcohol disorders	0.67 (0.58–0.78)	<0.001
Neurological disorders	0.81 (0.75–0.88)	<0.001
Psychiatric disorder	0.86 (0.80–0.94)	0.0004
Respiratory dysfunction	1.01 (0.92–1.12)	0.79
Vascular disease	0.77 (0.73–0.80)	<0.001

AMI-MM indicates acute myocardial infarction medically managed; CABG, coronary artery bypass grafting; CAD, coronary artery disease; COPD, chronic obstructive pulmonary disease; CR, cardiac rehabilitation; FFS, fee-for-service; HCC, hierarchical condition category; and PCI, percutaneous coronary intervention.

including 18 678 AMI-MM episodes, 41 986 PCI episodes, and 11 039 CABG episodes. A total of 20 613 (28.8%) of episodes had any CR enrollment within 90-days of hospital discharge. CR enrollment significantly varied by treatment strategy: CABG (53.8%), PCI (29.0%), AMI-MM (13.4%),  $P<0.001$ . Among all episodes, patients enrolled in CR were significantly different with respect to sex, age, prior spending, payer, admission status, and comorbidities (Table 1). Quarterly trends in CR enrollment for all CAD episodes of care, and by treatment strategy are displayed in Figure 1. There were significant differences in CR enrollment across payers, with commercial beneficiaries having the highest enrollment rates across all CAD events (35.8%),





**Figure 3.** Hospital variation in cardiac rehabilitation (CR) enrollment during coronary artery disease (CAD) episodes of care and by treatment strategy: medically managed acute myocardial infarction (AMI-MM), percutaneous coronary intervention (PCI), and coronary artery bypass grafting (CABG).

followed by Medicare Advantage (27.7%) and Medicare FFS (25.5%;  $P<0.001$ ; Figure 2). Significant differences in CR enrollment across payers was also found across treatment strategy (all  $P<0.001$ ), with commercial beneficiaries consistently having the highest CR enrollment rates.

**Predictors of CR Enrollment**

In all CAD episodes of care, CR enrollment was significantly lower among females versus males (adjusted odds ratio [aOR], 0.93 [95% CI, 0.90–0.97],  $P<0.001$ ), for Medicare FFS (aOR, 0.70 [95% CI, 0.67–0.73];  $P<0.001$ ) and Medicare Advantage (aOR, 0.70 [95% CI, 0.66–0.75];  $P<0.001$ ) versus commercial payers. Patients with several present comorbidities were also less likely to enroll in CR, including those with cerebrovascular disease, chronic kidney disease, chronic liver disease, chronic obstructive pulmonary disease, congestive heart failure, diabetes, drug/alcohol disorders, neurological disorders, psychiatric disorders, and vascular disease (Table 2). These patterns were similar across treatment strategies (Table III in the [Data Supplement](#)). Additionally, patients undergoing PCI with AMI versus without AMI were more likely to have enrolled in CR (aOR, 1.42 [95% CI, 1.39–1.49];  $P<0.001$ ), while CABG episodes with AMI versus without

AMI were less likely to have enrolled in CR (aOR, 0.76 [95% CI, 0.69–0.83];  $P<0.001$ ). These findings were similar when limiting the analysis to patients who were discharged home (Table IV in the [Data Supplement](#)).

**Hospital Variation in CR Enrollment**

Adjusting for patient factors, rates of CR enrollment varied from 9.8% to 51.6% across 33 hospitals following all CAD episodes of care (Figure 3). From the hierarchical logistic regression model, 9.3% of the variation was attributed to the hospital level, with a MOR of 1.73 (95% CI, 1.48–1.96; Table 3). In other words, the median relative odds in CR enrollment between 2 randomly chosen hospitals was 1.73.

Across treatment strategies, CR enrollment varied from 4.6% to 29.7% in AMI-MM episodes, from 7.5% to 58.5% in PCI episodes, and from 16.8% to 76.3% in CABG episodes (Figure 3). The ICC and MOR estimates of hospital variation were the greatest for PCI (ICC=10.7%, MOR=1.82), followed by AMI-MM (ICC=10.0%, MOR=1.78) and CABG (ICC=6.6%, MOR=1.58; Table 3). Within hospitals, CR enrollment by treatment strategy was highly correlated, with the strongest correlation between AMI-MM versus PCI ( $R^2=0.72$ ,  $P<0.001$ ), followed by PCI versus

**Table 3.** Hospital Variance Estimates for CR Enrollment During CAD Episodes of Care and by Treatment Strategy

Hospital variance estimates	All CAD events	AMI-MM	PCI	CABG
Risk-adjusted CR use, mean (range)	29.8% (9.8%–51.6%)	14.7% (4.6%–29.7%)	31.5 (7.5%–58.4%)	51.6 (16.8%–76.3%)
Hospital random effect variance (SE)	0.33 (0.09)	0.37 (0.10)	0.39 (0.10)	0.23 (0.07)
Intraclass correlation coefficient (95% CI)	9.1% (4.5%–13.3%)	10.1% (5.0%–14.7%)	10.6% (5.6%–15.2%)	6.5% (2.7%–10.0%)
Median odds ratio (95% CI)	1.73 (1.45–1.97)	1.78 (1.49–2.04)	1.81 (1.52–2.07)	1.58 (1.34–1.78)

AMI-MM indicates acute myocardial infarction medically managed; CABG, coronary artery bypass grafting; CAD, coronary artery disease; CR, cardiac rehabilitation, and PCI, percutaneous coronary intervention.




CABG ( $R^2=0.51$ ,  $P<0.001$ ) and AMI-MM versus CABG ( $R^2=0.46$ ,  $P<0.001$ ; Figure 4). We do note that hospitals with relatively low CR enrollment rates for AMI had higher rates of enrollment for PCI and CABG.

Table 4 shows CR enrollment, number of CR sessions attended, and the number of days to the first CR session during the 90-day CAD episodes of care by the tertile of risk-adjusted CR enrollment rate. CR enrollment increased significantly across tertiles: low=18.2%, moderate=27.1% and high=40.9% ( $P$ -trend<0.001). Among patients enrolled in CR, there were increasing trends in the mean number of CR sessions attended within 90 days of discharge (low=11.3 sessions, moderate=12.1 sessions, high=12.7 sessions,  $P$ -trend<0.001), and decreasing trend in the mean days to first session (low=44 days, moderate=40 days, high=34 days,  $P$ -trend<0.001). These patterns were similar across treatment strategies.

## DISCUSSION

In this analysis of 71 703 episodes of CAD at 33 hospitals in a statewide, multipayer claims registry, we report 4 major findings. First, there was over 5-fold variation in CR enrollment across hospitals during CAD episodes of care. Second, CR enrollment within a hospital was similar across CAD treatment strategies, suggesting hospital-specific, rather than treatment-specific, practice patterns. Third, the effect of payer and admitting hospital on CR enrollment was as large as or greater than patient demographic and clinical factors. Finally, patients in high CR enrollment hospitals had on average more CR sessions within 90 days of discharge and fewer days from discharge to first CR session.

Findings from this study have important implications for providers and policymakers seeking to improve CR enrollment. This study is among the first to highlight variation in hospital-level CR enrollment across multiple payers and CAD admission types. Importantly, prior studies have reported similar hospital-level variation and MORs for CR enrollment among Medicare beneficiaries and Veterans Affairs hospitals.<sup>19</sup> Moreover, a significant amount of the variation in CR enrollment was attributed to the hospital level, with similar enrollment rates across CAD patient cohorts receiving different treatments at the same hospital. Together, these findings suggest that hospitals may have specific practice patterns that may contribute to variation in CR enrollment. These findings differ from current literature, which demonstrates variation in CR enrollment at a regional level.<sup>7</sup> The underlying reasons for this variation may be a product of hospital-level practices, such as standardized referral protocols, affiliations with CR facilities, and physician endorsement, in addition to patient and local socioeconomic factors.<sup>20–23</sup> Nevertheless, several strategies have been shown to improve CR enrollment across hospital systems. The Million Hearts Initiative provides a



**Figure 4.** Within-hospital correlation in risk-adjusted cardiac rehabilitation (CR) enrollment rates during coronary artery disease episodes of care (medically managed acute myocardial infarction [AMI-MM], percutaneous coronary intervention [PCI], and coronary artery bypass grafting [CABG]).

framework for improving CR enrollment at the system level, which includes enhanced patient education, improving the process of enrolling patients into CR after

**Table 4.** Number of CR Claims and Days to First Claim Within 90 d of Discharge by Tertile of Risk-Adjusted CR Use

CR utilization	Tertile of risk-adjusted CR enrolment			P-trend
	Low	Moderate	High	
All CAD events (n=71 703)				
Patients, n	22 413	26 324	22 966	...
Any use, n (%)	4088 (18.2)	7131 (27.1)	9394 (40.9)	<0.0001
No. of claims, mean (SD)	11.3 (7.4)	12.1 (8.0)	12.7 (8.3)	<0.0001
Days to first claim, mean (SD)	44 (19)	40 (20)	34 (19)	<0.0001
Enrollment within 21 d, n (%)	440 (2.0)	1259 (4.8)	2774 (12.1)	<0.0001
AMI-MM (n=18678)				
Patients, n	5312	7491	5875	...
Any use, n (%)	347 (6.5)	843 (11.3)	1321 (22.5)	<0.0001
No. of claims, mean (SD)	11.6 (8.0)	11.0 (7.7)	12.0 (8.6)	0.029
Days to first claim, mean (SD)	42 (20)	42 (21)	35 (21)	<0.0001
Enrollment within 21 d, n (%)	52 (1.0)	159 (2.1)	451 (7.7)	<0.0001
PCI (n=41986)				
Patients, n	14 316	16 955	10 715	...
Any use, n (%)	2504 (17.5)	4819 (28.4)	4846 (45.2)	<0.0001
No. of claims, mean (SD)	11.8 (7.8)	12.5 (8.2)	13.1 (8.6)	<0.0001
Days to first claim, mean (SD)	40 (20)	38 (19)	30 (18)	<0.0001
Enrollment within 21 d, n (%)	438 (3.1)	1079 (6.4)	2044 (19.1)	<0.0001
CABG (n=11 039)				
Patients, n	2861	3851	4327	...
Any use, n (%)	1174 (41.0)	1970 (51.2)	2789 (64.5)	<0.0001
No. of claims, mean (SD)	10.4 (6.6)	11.4 (7.2)	12.6 (7.4)	<0.0001
Days to first claim, mean (SD)	53 (16)	46 (17)	41 (15)	<0.0001
Enrollment within 21 d, n (%)	11 (0.4)	77 (2.0)	162 (3.7)	<0.0001

AMI-MM indicates acute myocardial infarction medically managed; CABG, coronary artery bypass grafting; CAD, coronary artery disease; CR, cardiac rehabilitation; and PCI, percutaneous coronary intervention.

discharge, or tailoring CR programs to meet patient needs.<sup>24</sup> A systematic review of the literature showed that CR enrollment can be improved through both face-to-face delivery of CR with health care providers as well as offering unsupervised options.<sup>25</sup> Identifying and implementing interventions that are sensitive to the local resources of the center will increase the effectiveness of CR improvement efforts.

This study also adds to the growing body of literature demonstrating low rates of CR enrollment during CAD episodes of care. Over the 4 years in the study, there were only modest changes in CR enrollment, suggesting little progress has been made to improve enrollment. When examining CR enrollment across treatment strategies, our findings confirmed recent research demonstrating that medically managed AMI patients had lower rates of CR enrollment compared with those receiving revascularization procedures, such as PCI and CABG.<sup>7,26</sup> Enrollment rates in this study were also much lower than previously reported rates of referral to CR, highlighting the significant disparity between CR referral and enrollment. For example, reported rates

of referral to CR for PCI patients have exceeded 50% of patients,<sup>9,27</sup> yet we found less than a third of PCI recipients enrolled in a single session. Referral to CR is a quality measure endorsed by the National Quality Forum and recommended by the American College of Cardiology and American Heart Association.<sup>3</sup> While possibly easier to capture in real time, referral rates may be an insufficient proxy for enrollment.

Finally, we identify important patient-level factors associated with CR enrollment across CAD treatment strategies. This study confirms other reports that suggest age, female sex, and the presence of comorbidities are significant predictors of CR enrollment.<sup>7,15,26,28–30</sup> The findings also confirm that patients undergoing PCI with AMI are more likely to enroll in CR after discharge, while CABG patients with AMI are less likely to enroll in CR.<sup>7,8</sup> Moreover, we add to the literature on patient determinants of CR enrollment by highlighting differences across payers. Specifically, we found that CR enrollment was the greatest among the commercially insured population compared with Medicare FFS or Medicare Advantage, independent of the treatment

strategy chosen. We do note that differences across payers may be partly explained by socioeconomic factors not included in our data. Prior studies have shown socioeconomic indicators to be linked to CR referral and enrollment.<sup>7,31,32</sup> To improve CR enrollment, more work is needed to understand barriers related to insurance coverage and cost-sharing, which could inform public and private payer benefit designs.

These findings should be interpreted with several limitations in mind. First, data used in this study are limited to CAD episodes of care in Michigan and to beneficiaries with Medicare FFS and BCBSM/BCN commercial and Medicare Advantage plans, which may not be generalizable to all CAD episodes of care. However, this study expands upon existing literature that often focus solely on Medicare FFS beneficiaries by including beneficiaries from other payers.<sup>7,26,32</sup> Second, we only captured CR enrollment within 90 days of discharge, which likely underestimates overall enrollment rates and the number of CR sessions attended. Thus, longer time to enrollment may result in fewer CR sessions within the 90-day period. Third, there are limitations to claims data vis-à-vis missing data. Claims data lack clinically relevant and validated measures of disease severity, clinical status, and procedural information that might be important determinants of CR enrollment. Information may also be missing if it is irrelevant to medical billing practices, which may differ across hospitals. Future work should incorporate clinical registries into claims data to address these missing data issues. Fourth, administrative data also lack information on CR referral, which is an important predictor of enrollment and may explain variation in enrollment across hospitals.<sup>8,15</sup> Fifth, we do not include clustering or analysis of variation at the physician level. Grace et al<sup>21</sup> identified that physician endorsement of CR was a significant determinant of enrollment, which may be a contributing factor to variation across hospitals that was not available in our data.<sup>33</sup> Sixth, we were not able to separate out individuals with type II non-ST elevated myocardial infarction, which have historically lower rates of CR enrollment.<sup>34</sup> Finally, episodes of care in this study do not include patients with stable CAD, heart failure, valve repair/replacement, or heart transplant, who are also eligible for CR enrollment.

## CONCLUSIONS

Enrollment in CR during CAD episodes of care varies from 10% to 50% across hospitals and treatment strategy. While several patient demographics, clinical, and insurance factors predicted eventual enrollment in CR, 10% of the variation in enrollment was attributed to the hospital-level. Future work should seek to identify and implement hospital-specific practices associated with higher CR enrollment and address nonhospital

barriers to enrollment to facilitate national efforts supported by federal agencies and payers.

## ARTICLE INFORMATION

Received August 25, 2020; accepted November 23, 2020.

This manuscript was sent to Dennis T. Ko, MD, MSc, Senior Guest Editor, for review by expert referees, editorial decision, and final disposition.

The Data Supplement is available at <https://www.ahajournals.org/doi/suppl/10.1161/CIRCOUTCOMES.120.007144>.

## Correspondence

Michael P. Thompson, PhD, 5331K Frankel Cardiovascular Center, 1500 E Medical Center Dr, SPC 5864, Ann Arbor, MI 48109. Email [mthomps@med.umich.edu](mailto:mthomps@med.umich.edu)

## Affiliations

Department of Cardiac Surgery (M.P.T., H.H., D.S.L., R.L.P.) and Division of Cardiovascular Medicine, Department of Internal Medicine (H.S.G., D.S.), Michigan Medicine, Ann Arbor MI. Michigan Value Collaborative, University of Michigan, Ann Arbor (M.P.T., J.M.Y., J.D.S.). Department of Cardiac Surgery, Bronson Methodist Hospital, Kalamazoo, MI (A.D.). Michigan Society of Thoracic and Cardiovascular Surgeons Quality Collaborative, Ann Arbor, MI (D.S.L., R.L.P.). Division of Cardiovascular Medicine, Henry Ford Health System, Detroit, MI (S.J.K.). Blue Cross Blue Shield of Michigan Cardiovascular Consortium, (BMC2), Ann Arbor, MI (H.S.G., D.S.).

## Sources of Funding

None.

## Disclosures

Dr Thompson receives partial salary support from Blue Cross Blue Shield of Michigan. Dr Sukul receives partial salary support from Blue Cross Blue Shield of Michigan. Dr Sukul is a member of the Government Relations Committee for the Society for Cardiovascular Angiography & Interventions, which had no role in the work herein. Dr Likosky is a consultant to the American Society of Extracorporeal Technology and receives funding from the Agency for Healthcare Research and Quality (R01HS026003) and the National Institutes of Health (HL146619-01A1). Dr Gurm receives research support from Blue Cross and Blue Shield of Michigan, the National Institutes of Health Center for Accelerated Innovations, and Michigan Translational Research and Commercialization for Life Sciences Innovation Hub. He is the co-founder of, owns equity in, and is a consultant to Amplitude Vascular Systems. He also owns equity in Jiaxing Bossh Medical Technology Partnership and is a consultant for Osprey Medical. He is the chair of the Clinical Events Committee for the Protection Against Emboli During Carotid Artery Stenting Using the Neuroguard IEP System trial sponsored by Contego Medical. The other authors report no conflicts.

## Supplemental Material

Tables I–IV

## REFERENCES

1. Heran BS, Chen JM, Ebrahim S, Moxham T, Oldridge N, Rees K, Thompson DR, Taylor RS. Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database Syst Rev*. 2011;CD001800.
2. Anderson L, Oldridge N, Thompson DR, Zwisler AD, Rees K, Martin N, Taylor RS. Exercise-based cardiac rehabilitation for coronary heart disease: Cochrane systematic review and meta-analysis. *J Am Coll Cardiol*. 2016;67:1–12. doi: 10.1016/j.jacc.2015.10.044
3. Thomas RJ, Balady G, Banka G, Beckie TM, Chiu J, Gokak S, Ho PM, Keteyian SJ, King M, Lui K, Pack Q, Sanderson BK, Wang TY. 2018 ACC/AHA clinical performance and quality measures for cardiac rehabilitation: a report of the American College of Cardiology/American Heart Association Task Force on Performance Measures. *Circ Cardiovasc Qual Outcomes*. 2018;11:e000037. doi: 10.1161/HCQ.0000000000000037

4. Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C, Catapano AL, Cooney MT, Corrà U, Cosyns B, Deaton C, et al; ESC Scientific Document Group. 2016 European Guidelines on cardiovascular disease prevention in clinical practice: the Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts) Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J*. 2016;37:2315–2381. doi: 10.1093/eurheartj/ehw106
5. Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr, Colvin MM, Drazner MH, Filippatos GS, Fonarow GC, Givertz MM, et al. 2017 ACC/AHA/HFSA focused update of the 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation*. 2017;136:e137–e161. doi: 10.1161/CIR.0000000000000509
6. Hirsch AT, Haskal ZJ, Hertzner NR, Bakal CW, Creager MA, Halperin JL, Hiratzka LF, Murphy WR, Olin JW, Puschett JB, et al; American Association for Vascular Surgery; Society for Vascular Surgery; Society for Cardiovascular Angiography and Interventions; Society for Vascular Medicine and Biology; Society of Interventional Radiology; ACC/AHA Task Force on Practice Guidelines Writing Committee to Develop Guidelines for the Management of Patients With Peripheral Arterial Disease; American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; TransAtlantic Inter-Society Consensus; Vascular Disease Foundation. ACC/AHA 2005 practice guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic): a collaborative report from the American Association for Vascular Surgery/Society for Vascular Surgery, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients With Peripheral Arterial Disease): endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; TransAtlantic Inter-Society Consensus; and Vascular Disease Foundation. *Circulation*. 2006;113:e463–e654. doi: 10.1161/CIRCULATIONAHA.106.174526
7. Ritchey MD, Maresh S, McNeely J, Shaffer T, Jackson SL, Keteyian SJ, Brawner CA, Whooley MA, Chang T, Stolp H, et al. Tracking cardiac rehabilitation participation and completion among medicare beneficiaries to inform the efforts of a national initiative. *Circ Cardiovasc Qual Outcomes*. 2020;13:e005902. doi: 10.1161/CIRCOUTCOMES.119.005902
8. Sukul D, Seth M, Barnes GD, Dupree JM, Syrjamaki JD, Dixon SR, Maddler RD, Lee D, Gurm HS. Cardiac rehabilitation use after percutaneous coronary intervention. *J Am Coll Cardiol*. 2019;73:3148–3152. doi: 10.1016/j.jacc.2019.03.515
9. Beatty AL, Bradley SM, Maynard C, McCabe JM. Referral to cardiac rehabilitation after percutaneous coronary intervention, coronary artery bypass surgery, and valve surgery: data from the clinical outcomes assessment program. *Circ Cardiovasc Qual Outcomes*. 2017;10:1–8.
10. Peters AE, Keeley EC. Trends and predictors of participation in cardiac rehabilitation following acute myocardial infarction: data from the behavioral risk factor surveillance system. *J Am Heart Assoc*. 2017;7:1–6.
11. Marzolini S, Blanchard C, Alter DA, Grace SL, Oh PI. Delays in referral and enrollment are associated with mitigated benefits of cardiac rehabilitation after coronary artery bypass surgery. *Circ Cardiovasc Qual Outcomes*. 2015;8:608–620.
12. Dunlay SM, Witt BJ, Allison TG, Hayes SN, Weston SA, Koepsell E, Roger VL. Barriers to participation in cardiac rehabilitation. *Am Heart J*. 2009;158:852–859. doi: 10.1016/j.ahj.2009.08.010
13. Niebauer Josef. Is there a role for cardiac rehabilitation after coronary artery bypass grafting? *Circulation*. 2016;133:2529–2537.
14. Ellimootil C, Syrjamaki JD, Voit B, Guduguntla V, Miller DC, Dupree JM. Validation of a claims-based algorithm to characterize episodes of care. *Am J Manag Care*. 2017;23:e382–e386.
15. Doll JA, Hellkamp A, Ho PM, Kontos MC, Whooley MA, Peterson ED, Wang TY. Participation in cardiac rehabilitation programs among older patients after acute myocardial infarction. *JAMA Intern Med*. 2015;175:1700–1702. doi: 10.1001/jamainternmed.2015.3819
16. Ellis RP, Pope GC, Iezzoni L, Ayanian JZ, Bates DW, Burstin H, Ash AS. Diagnosis-based risk adjustment for Medicare capitation payments. *Health Care Financ Rev*. 1996;17:101–128.
17. Pope GC, Kautter J, Ellis RP, Ash AS, Ayanian JZ, Iezzoni LI, Ingber MJ, Levy JM, Robst J. Risk adjustment of Medicare capitation payments using the CMS-HCC model. *Health Care Financ Rev*. 2004;25:119–141.
18. Merlo J, Chaix B, Ohlsson H, Beckman A, Johnell K, Hjerpe P, Råstam L, Larsen K. A brief conceptual tutorial of multilevel analysis in social epidemiology: using measures of clustering in multilevel logistic regression to investigate contextual phenomena. *J Epidemiol Community Health*. 2006;60:290–297. doi: 10.1136/jech.2004.029454
19. Beatty AL, Doll JA, Schopfer DW, Maynard C, Plomondon ME, Shen H, Whooley MA. Cardiac rehabilitation participation and mortality after percutaneous coronary intervention: insights from the veterans affairs clinical assessment, reporting, and tracking program. *J Am Heart Assoc*. 2018;7:e010010. doi: 10.1161/JAHA.118.010010
20. Dalal HM, Doherty P, Taylor RS. Cardiac rehabilitation. *BMJ*. 2015;351:h5000. doi: 10.1136/bmj.h5000
21. Grace SL, Gravely-Witte S, Brual J, Monette G, Suskin N, Higginson L, Alter DA, Stewart DE. Contribution of patient and physician factors to cardiac rehabilitation enrollment: a prospective multilevel study. *Eur J Cardiovasc Prev Rehabil*. 2008;15:548–556. doi: 10.1097/HJR.0b013e328305df05
22. Shanmugasagaram S, Oh P, Reid RD, McCumber T, Grace SL. Cardiac rehabilitation barriers by rurality and socioeconomic status: a cross-sectional study. *Int J Equity Health*. 2013;12:72. doi: 10.1186/1475-9276-12-72
23. Grace SL, Evindar A, Abramson BL, Stewart DE. Physician management preferences for cardiac patients: factors affecting referral to cardiac rehabilitation. *Can J Cardiol*. 2004;20:1101–1107.
24. CDC. Million Hearts® Cardiac Rehabilitation Change Package [Internet]. Centers for Disease Control and Prevention. 2019. <https://millionhearts.hhs.gov/tools-protocols/action-guides/cardiac-change-package/index.html>. Accessed December 23, 2019.
25. Pio CSA, Chaves G, Davies P, Taylor R, Grace S. Interventions to promote patient utilization of cardiac rehabilitation: cochrane systematic review and meta-analysis. *J Clin Med Res*. 2019;8:1–19.
26. Suaya JA, Shepard DS, Normand SL, ADES PA, Protas J, Stason WB. Use of cardiac rehabilitation by medicare beneficiaries after myocardial infarction or coronary bypass surgery. *Circulation*. 2007;116:1653–1662. doi: 10.1161/CIRCULATIONAHA.107.701466
27. Aragam KG, Dai D, Neely ML, Bhatt DL, Roe MT, Rumsfeld JS, Gurm HS. Gaps in referral to cardiac rehabilitation of patients undergoing percutaneous coronary intervention in the United States. *J Am Coll Cardiol*. 2015;65:2079–2088. doi: 10.1016/j.jacc.2015.02.063
28. Pack QR, Goel K, Lahr BD, Greason KL, Squires RW, Lopez-Jimenez F, Zhang Z, Thomas RJ. Participation in cardiac rehabilitation and survival after coronary artery bypass graft surgery. *Circulation*. 2013;128:590–597.
29. Menezes AR, Lavie CJ, Milani RV, Arena RA, Church TS. Cardiac rehabilitation and exercise therapy in the elderly: should we invest in the aged? *J Geriatr Cardiol*. 2012;9:68–75. doi: 10.3724/SP.J.1263.2012.00068
30. Grace SL, Shanmugasagaram S, Gravely-Witte S, Brual J, Suskin N, Stewart DE. Barriers to cardiac rehabilitation: DOES AGE MAKE A DIFFERENCE? *J Cardiopulm Rehabil Prev*. 2009;29:183–187. doi: 10.1097/HCR.0b013e3181a3333c
31. Castellanos LR, Viramontes O, Bains NK, Zepeda IA. Disparities in cardiac rehabilitation among individuals from racial and ethnic groups and rural communities—a systematic review. *J Racial Ethn Health Disparities*. 2019;6:1–11. doi: 10.1007/s40615-018-0478-x
32. Patel DK, Duncan MS, Shah AS, Lindman BR, Greevy RA Jr, Savage PD, Whooley MA, Matheny ME, Freiberg MS, Bachmann JM. Association of cardiac rehabilitation with decreased hospitalization and mortality risk after cardiac valve surgery. *JAMA Cardiol*. 2019;4:1250–1259. doi: 10.1001/jamacardio.2019.4032
33. Ghisi GL, Polyzotis P, Oh P, Pakosh M, Grace SL. Physician factors affecting cardiac rehabilitation referral and patient enrollment: a systematic review. *Clin Cardiol*. 2013;36:323–335. doi: 10.1002/clc.22126
34. McCarthy CP, Murphy S, Cohen JA, Rehman S, Jones-O'Connor M, Olshan DS, Singh A, Vaduganathan M, Januzzi JL Jr, Wasfy JH. Underutilization of cardiac rehabilitation for type 2 myocardial infarction. *J Am Coll Cardiol*. 2019;73:2005–2007. doi: 10.1016/j.jacc.2019.01.032