Henry Ford Health Henry Ford Health Scholarly Commons

Cardiology Articles

Cardiology/Cardiovascular Research

9-1-2021

Fitness and Mortality Among Persons 70 Years and Older Across the Spectrum of Cardiovascular Disease Risk Factor Burden: The FIT Project

Seamus P. Whelton

Paul A. McAuley

Zeina Dardari

Olusola A. Orimoloye

Erin D. Michos

See next page for additional authors

Follow this and additional works at: https://scholarlycommons.henryford.com/cardiology_articles

Recommended Citation

Whelton SP, McAuley PA, Dardari Z, Orimoloye OA, Michos ED, Brawner CA, Ehrman JK, Keteyian SJ, Blaha MJ, and Al-Mallah MH. Fitness and Mortality Among Persons 70 Years and Older Across the Spectrum of Cardiovascular Disease Risk Factor Burden: The FIT Project. Mayo Clin Proc 2021.

This Article is brought to you for free and open access by the Cardiology/Cardiovascular Research at Henry Ford Health Scholarly Commons. It has been accepted for inclusion in Cardiology Articles by an authorized administrator of Henry Ford Health Scholarly Commons.

Authors

Seamus P. Whelton, Paul A. McAuley, Zeina Dardari, Olusola A. Orimoloye, Erin D. Michos, Clinton A. Brawner, Jonathan K. Ehrman, Steven J. Keteyian, Michael J. Blaha, and Mouaz H. Al-Mallah



Check for updates

Fitness and Mortality Among Persons 70 Years and Older Across the Spectrum of Cardiovascular Disease Risk Factor Burden: The FIT Project

Seamus P. Whelton, MD, MPH; Paul A. McAuley, PhD; Zeina Dardari, MS; Olusola A. Orimoloye, MD, MPH; Erin D. Michos, MD, MHS; Clinton A. Brawner, PhD; Jonathan K. Ehrman, PhD; Steven J. Keteyian, PhD; Michael J. Blaha, MD, MPH; and Mouaz H. Al-Mallah, MD, MSc

Abstract

Objective: To determine whether fitness could improve mortality risk stratification among older adults compared with cardiovascular disease (CVD) risk factors.

Methods: We examined 6509 patients 70 years of age and older without CVD from the Henry Ford ExercIse Testing Project (FIT Project) cohort. Patients performed a physician-referred treadmill stress test between 1991 and 2009. Traditional categorical CVD risk factors (hypertension, hyperlipidemia, diabetes, and smoking) were summed from 0 to 3 or more. Fitness was grouped as low, moderate, and high (<6, 6 to 9.9, and \geq 10 metabolic equivalents of task). All-cause mortality was ascertained through US Social Security Death Master files. We calculated age-adjusted mortality rates, multivariable adjusted Cox proportional hazards, and Kaplan-Meier survival models.

Results: Patients had a mean age of 75 ± 4 years, and 3385 (52%) were women; during a mean followup of 9.4 years, there were 2526 deaths. A higher fitness level (*P*<.001), not lower CVD risk factor burden (*P*=.31), was associated with longer survival. The age-adjusted mortality rate per 1000 personyears was 56.7 for patients with low fitness and 0 risk factors compared with 24.9 for high fitness and 3 or more risk factors. Among patients with 3 or more risk factors, the adjusted mortality hazard was 0.68 (95% CI, 0.61 to 0.76) for moderate and 0.51 (95% CI, 0.44 to 0.60) for high fitness compared with the least fit.

Conclusion: Among persons aged 70 years and older, there was no significant difference in survival of patients with 0 vs 3 or more risk factors, but a higher fitness level identified older persons with good long-term survival regardless of CVD risk factor burden.

© 2021 Mayo Foundation for Medical Education and Research
Mayo Clin Proc. 2021;96(9):2376-2385

0

From the Johns Hopkins Ciccarone Center for the Prevention of Cardiovascular Disease, Johns Hopkins School of Medicine, Baltimore, MD (S.P.W., Z.D., O.A.O., E.D.M., M.J.B.); the Department of Health, Physical Education, and Sport Studies, Winston-Salem

Affiliations continued at the end of this article.

R isk stratification in persons 70 years of age and older presents a significant challenge because with increasing age, there is both an increased prevalence of cardiovascular disease (CVD) risk factors and a reduction in their predictive strength.¹⁻³ There is also a higher absolute rate of both CVD and non-CVD morbidity and mortality in older age, which contributes to an increase in competing risks.⁴⁻⁶ Accordingly, for older persons, age

functions as the dominant predictor in most of the commonly used risk prediction models, leading to nearly all individuals 70 years and older being classified as high risk, whether they have 0 or multiple CVD risk factors.⁷⁻⁹ Whereas CVD risk factors are most strongly associated with CVD outcomes, they are notably also associated with an increased risk for total mortality, an especially important outcome to consider among older persons.¹⁰⁻¹⁴ Therefore, additional variables are needed to improve risk stratification among older individuals.

Although fitness is not included in commonly used risk scores, it is one of the strongest predictors for the risk of both CVD and total mortality in the general population, and among middle-aged persons, treadmill exercise testing abnormalities significantly improve risk stratification beyond traditional CVD risk factors.15-22 However, fitness decreases with age, and its utility for risk stratification is less well defined among older individuals.²³ It is uncertain whether older individuals have differences in their mean fitness level across a low vs high CVD risk factor burden. In addition, it is unknown whether the association between fitness and mortality differs by risk factor burden. Therefore, we sought to determine the association between fitness and mortality among persons 70 years of age and older across the spectrum of CVD risk factor burden.

METHODS

The Henry Ford ExercIse Testing Project (FIT Project) is composed of 69,885 patients who performed a physician-referred exercise treadmill stress test between January 2, 1991, and May 27, 2009, at the Henry Ford Hospital (Detroit, Michigan) and affiliated medical centers. A detailed description of the study design, methods, follow-up, and mortality ascertainment has been previously reported.²⁴ For this analysis, we excluded patients who were younger than 70 years (n=60,488),those with prior CVD (n=2776), and those missing data on fitness (n=112), for a total of 6509 persons 70 years of age and older who were included in this study.

Treadmill stress testing was performed using a standard Bruce protocol per the American Heart Association/American College of Cardiology guideline recommended methodology.^{25,26} The test was stopped if the patient had exercise-limiting symptoms as determined by the supervising clinician (clinical exercise physiologist or nurse) or if the patient requested that the test be stopped. The test was also stopped if the patient had an abnormal blood pressure response (eg, a decrease in systolic blood pressure of >10 mm Hg despite an increased workload, systolic blood pressure >250 mm Hg, or diastolic blood pressure >115 mm Hg), significant ST-segment changes, clinically significant arrhythmia, or exerciselimiting chest pain. The patients' peak exercise workload (peak treadmill speed and peak incline) was used to calculate their maximal exercise capacity (eg, fitness), which was reported as metabolic equivalents of task (METs). Fitness was categorized as low (<6 METs), moderate (6-9 METs), and high (≥ 10 METs).²⁷ We chose 10 METs and higher as the high-fitness group because older persons have a lower mean fitness level and few patients in this study achieved 12 METs or more (n=74). We used physician referral information to group the stress test indications into common categories, such as chest pain and dyspnea.²⁴

Immediately preceding the treadmill stress test, a trained nurse or clinical exercise physiologist obtained a medical history including the patient's demographic characteristics, CVD risk factors, current smoking status, current medication use, and past medical history. This self-reported information was supplemented by a retrospective search of the administrative databases or pharmacy claims files for patients who were enrolled in the Henry Ford Health System integrated health plan to augment the information on medication use and past medical history. For an electronic medical record (EMR) diagnosis to be coded as such in our database, it must have been present on at least 3 separate clinical encounters.²⁴ We used these data combining patient self-report, use of a disease-specific medication, or an EMR database-verified diagnosis to classify hypertension, dyslipidemia, diabetes, and current smoking as present or absent.

The reported resting systolic and diastolic blood pressures were measured immediately before the stress test, but these values were not used to define hypertension for this study because a diagnosis of hypertension must be based on 2 separate measurements

on at least 2 separate visits, patients may have had higher than typical blood pressures on the day of their stress test because of withholding of medications for the stress test, and it was not recorded whether the blood pressure readings were measured with the patient in the standing or supine position. A family history of coronary artery disease was defined as the presence of a firstdegree relative with a prior clinical coronary artery disease event. For patients 80 years and older, an age of 79 years was used to calculate the pooled cohort equations (PCEs) as recommended by the American College of Cardiology/American Heart Association because the PCEs were developed using individuals aged 40 to 79 years. We performed a retrospective search of the EMR and associated laboratory databases to identify laboratory values obtained within 90 days of the stress test. There were 614 participants missing laboratory data for total cholesterol concentration, but EMR data for a diagnosis of hyperlipidemia or lipidlowering medication use were available for these patients. All-cause mortality was obtained by linkage to the Social Security Death Master File through April 2013.²⁴

Each patient's traditional CVD risk factor (hypertension, dyslipidemia, diabetes, and smoking) burden was summed from 0 to 3 or more, and the proportion of individuals within each risk factor group was stratified by fitness categories. The age-adjusted mortality rate was calculated per 1000 personyears of follow-up stratified by METs and fitness groups. Kaplan-Meier survival analysis was performed on the basis of the number of CVD risk factors and fitness categories.²⁸ The primary outcome was allcause mortality. Follow-up was calculated as the days between the exercise stress test date and the date of death or the end date for death matching (April 30, 2013). In addition, we performed multivariable adjusted Cox proportional hazards modeling to examine the risk of mortality based on the number of traditional CVD risk factors and fitness.²⁹ In our adjusted analyses, model 1 included age, sex, race, and reason for the stress test; model 2 additionally included

resting systolic and diastolic blood pressure, total cholesterol concentration, high-density lipoprotein cholesterol concentration, use of hypertension medication, use of lipidlowering medication, diabetes, and smoking status. We performed sensitivity analyses excluding patients with a diagnosis of cancer at baseline (n=777; defined as an EMR-listed diagnosis), stratified by sex, and excluding patients who died within the first year of follow-up (n=107).

RESULTS

Patients ranged in age from 70 to 96 years, with a mean age of 75 years (SD 4.2 years); 3385 (52%) were women, and 4603 (70.7%) were White (Table 1). Medication treatment for the risk factors occurred among 4269 (79%) patients with hypertension, 1766 (54%) patients with dyslipidemia, and 714 (43%) patients with diabetes (71 (10%) of whom were prescribed insulin). Patients who reported current smoking received smoking cessation advice per their referring physician's standard practice. The proportion of patients with CVD risk factors was lower with higher fitness levels (Supplemental Table 1, available online at http://www.mayoclinicproceedings.org). There were 2526 deaths during a mean follow-up of 9.4 years. The mean 10-year PCE risk score was 35.3%, and only 0.2% (n=10) had a 10-year PCE score below 7.5%. The CVD risk factor burden was similar for patients across fitness levels; 6125 (94.1%) patients had at least 1 CVD risk factor, whereas 1868 (28.7%) patients had 3 or more CVD risk factors (Figure 1). Regardless of the number of CVD risk factors, individuals in the low-fitness group had the highest age-adjusted mortality rate (56.7 per 1000 person-years), whereas patients in the highest fitness group had the lowest age-adjusted mortality rate (25.1 per 1000 person-years). In addition, within each fitness group, the age-adjusted mortality rate did not increase with a higher CVD risk factor burden (Figure 2). During 15 years of follow-up, there was no significant difference in survival for patients with 0 CVD risk factors compared with those

TABLE 1. Participant Characteristics Stratified by Number of Risk Factors ^{a,b,c}									
No. of risk factors	0 (n=384)	l (n=1743)	2 (n=2512)	≥3 (n=1870)	P for trend				
Age (y)	75.6 (4.3)	75.7 (4.5)	75.5 (4.2)	75.0 (3.8)	<.01				
Women	182 (47.4)	906 (52.0)	1392 (55.4)	905 (48.4)	.37				
Ethnicity									
White	303 (78.9)	1249 (71.7)	1781 (70.9)	1270 (67.9)	<.001				
African American	61 (15.9)	423 (24.3)	634 (25.2)	(27.8)	<.001				
Systolic blood pressure (mm Hg)	138.0 (19.5)	141.2 (19.7)	142.6 (19.7)	4 .7 (9.)	.01				
Diastolic blood pressure (mm Hg)	79.4 (10.1)	90.7 (10.3)	80.7 (10.1)	79.6 (10.0)	.17				
Hypertension	0 (0)	1253 (71.9)	2304 (91.7)	1849 (98.9)	<.001				
Hypertension medication	0 (0)	971 (55.7)	1792 (71.3)	1506 (80.5)	<.001				
Total cholesterol (mg/dL)	204.9 (39.7)	203.1 (39.1)	206.5 (41.4)	198.2 (42.2)	<.001				
HDL-C (mg/dL)	56.3 (17.5)	54.8 (16.9)	53.8 (15.7)	50.5 (15.0)	<.001				
Dyslipidemia	0 (0)	217 (12.5)	1415 (56.3)	1636 (87.5)	<.001				
Statin	0 (0)	72 (4.1)	696 (27.7)	998 (53.4)	<.001				
Diabetes	0 (0)	49 (2.8)	470 (18.7)	1138 (60.9)	<.001				
Current smoking	0 (0)	224 (12.9)	835 (33.2)	1288 (68.9)	<.001				
Reason for stress test									
Chest pain	162 (42.2)	761 (43.7)	1052 (41.9)	716 (38.3)	<.001				
Shortness of breath	37 (9.6)	189 (10.8)	318 (12.6)	252 (13.5)	<.001				
Rule out ischemia	59 (15.4)	227 (13.0)	334 (13.3)	254 (13.6)	.94				
Other	25 (32.8)	116 (32.5)	169 (32.2)	126 (34.6)	.47				
10-year ASCVD risk ^d	24.7 (12.2)	28.9 (13.6)	34.2 (15.3)	44.7 (15.9)	<.001				
METs achieved	6.4 (2.5)	6.2 (2.5)	6.2 (2.4)	6.3 (2.2)	.48				
Achieving >10 METS	82 (21.4)	301 (17.2)	425 (16.7)	265 (14.1)	<.001				

^aASCVD, atherosclerotic cardiovascular disease; HDL-C, high-density lipoprotein cholesterol; METs, metabolic equivalents of task.

^bTo convert cholesterol values to mmol/L, multiply by 0.0259.

^cCategorical variables are presented as number (%). Continuous variables are presented as mean (standard deviation).

^dASCVD risk estimated using the 2013 American Heart Association/American College of Cardiology pooled cohort equations.

with 3 or more CVD risk factors (P=.31; Figure 3A). However, higher fitness levels were associated with significantly longer survival, and during 15 years of follow-up, 669 (62.3%) patients in the highest fitness group were still alive compared with 1023 (35.4%) patients in the lowest fitness group (P<.001; Figure 3B).

On multivariable adjusted Cox proportional hazards modeling (Table 2), there was no significant difference in the hazard of death for patients with 3 or more CVD risk factors compared with those with 0 CVD risk factors (hazard ratio [HR], 1.18; 95% CI, 0.97 to 1.43; P=.10). Compared with patients in the lowest fitness group, those with moderate fitness had a 37% lower hazard of death (HR, 0.63; 95% CI, 0.57 to 0.68), and those with high fitness had a 56% lower hazard of death (HR, 0.44; 95%, CI, 0.38 to 0.50). Within each CVD risk factor group, there was approximately a 35% reduction in hazard of death for those with moderate fitness and a more than 50% reduction in hazard of death for those with high fitness compared with patients in the lowest fitness group (Supplemental Table 2, available online http://www. at mayoclinicproceedings.org). In addition, within each CVD risk factor group, there was approximately a 15% lower hazard of death for every 1 MET higher fitness achieved (P<.001). In sensitivity analyses, we found no significant difference in the results after excluding patients with a diagnosis of cancer at baseline, between women and men, or after excluding patients who died within the first year of follow-up.

Mayo Clin Proc. September 2021;96(9):2376-2385 https://doi.org/10.1016/j.mayocp.2020.12.039 www.mayoclinicproceedings.org



DISCUSSION

Among patients aged 70 years and older who were free of known CVD, we found no significant difference in long-term survival by the number of traditional CVD risk factors, but a higher fitness level was strongly associated with a considerably lower mortality rate across the spectrum of CVD risk factor burden. These results highlight the utility of fitness as a risk stratification tool among older individuals, a group in whom risk stratification by traditional risk factors is especially challenging. In addition, our findings add further evidence supporting the importance of fitness as a leading modifiable risk factor among older persons.^{30,31}

CVD risk stratification based on traditional CVD risk factors is likely to be the most commonly performed type of risk assessment in routine clinical practice. It may parenthetically also serve as a marker of overall long-term prognosis, particularly among older individuals, in whom CVD is the leading cause of death. Commonly used risk equations were derived using data from predominantly middle-aged individuals and are applied to older individuals, but this pragmatic approach overlooks meaningful physiologic differences between younger, middle-aged, and older persons.32-34 For instance, whereas age is heavily weighted in the PCE and serves as a surrogate for CVD risk factor duration, the relative strength of the association between CVD risk factors and outcomes significantly declines with increasing age.³⁵⁻³⁷ This is largely attributable to an increased prevalence of CVD risk factors with increasing age, consistent with our finding that 94% of individuals in this study had at least 1 CVD risk factor. In addition, higher absolute event rates for both CVD and non-CVD outcomes lead to increased competing risks.^{2,38} Therefore, variables that provide a measure of physiologic age rather than of chronologic age are imperative for accurate risk stratification among older persons.^{39,40}

Whereas the absence of CVD risk factors among older patients is typically considered a healthy phenotype, nearly no men older than 65 years or women older than 70 years are deemed low risk (10-year risk, <7.5%) by the PCE, even in the absence of traditional CVD risk factors.⁴¹ Consequently, this limited ability to identify healthy older individuals is a weakness of commonly used risk scores that are based on traditional CVD risk factors.⁴⁰ In contrast, our results

demonstrate that higher fitness identifies older patients with good long-term survival regardless of their CVD risk factor burden. Indeed, we found that patients with a high fitness level and 3 or more CVD risk factors had a nearly 60% lower mortality rate compared with patients without any CVD risk factors but a low fitness level.

The ability of higher fitness to identify older persons with good long-term survival is an equally important strength compared with the identification of older persons with poor long-term survival based on a low fitness level. This robustness to identify both low- and highrisk older persons is attributable to the dependence of fitness on the global cardiovascular health of an individual. Accordingly, fitness indirectly integrates measures of an individual's myocardial function, coronary and peripheral atherosclerotic burden, vascular endothelial function, and pulmonary function in addition to other important risk predictors, such as large-muscle strength and frailty.⁴²⁻⁴⁶ The measurement of traditional CVD risk factors does not cohesively integrate these physiologic measures, which is at least partly responsible for the near absence of older individuals who are classified as low risk by commonly used risk prediction models.

Previous attempts to improve risk stratification in the elderly have shown only a limited improvement compared with traditional risk scores. For example, Koller et al47 used data from the Cardiovascular Health Study and Rotterdam Study to develop a coronary heart disease risk prediction model in persons 65 years of age and older. Unfortunately, the risk score derived from the elderly population showed only a small improvement in risk prediction even after the addition of novel risk factors, such as ankle-brachial index, carotid intimal thickness, and C-reactive protein level. Vliegenthart et al,48 also using data from the Rotterdam Study, investigated the utility of coronary artery calcium (CAC) scoring for risk prediction among individuals aged 70 years and older. Whereas they found a highly significant relative association between CAC and adverse outcomes, the addition of CAC to a multivariate model showed an



factors.

increase in the area under the curve of only 0.013 for CVD and 0.006 for mortality.

Although traditional CVD risk factors are most strongly associated with CVD mortality, this study's primary outcome of allcause mortality remains an important and relevant outcome to examine. A better understanding of one's long-term mortality risk may be important for a patient's selfawareness of overall health, identifying the patient's priorities for health care treatment and quality of life along with identifying long-term goals of care.49 Indeed, the National Institute for Health and Care Excellence highlighted the need to develop new methods for robust mortality risk stratification among patients with multiple chronic diseases, which is common among older persons.⁵⁰ Mortality risk prediction equations such as QMortality, Hospital-Patient One-Year Mortality Risk, and the Johns Hopkins Aggregated Diagnostic Groups predict only short-term mortality (1-2 years) and contain

Mayo Clin Proc. September 2021;96(9):2376-2385 https://doi.org/10.1016/j.mayocp.2020.12.039 www.mayoclinicproceedings.org



between 20 and 57 variables.^{49,51,52} Therefore, they have a relatively narrow utility in routine patient care.

The proportion of deaths due to CVD also increases sharply above the age of 70 years and accounts for approximately one-third of all deaths in this age group (n=508,804 deaths among persons \geq 75 years old in the United States in 2016).⁵³ Furthermore, the CVD risk factors included in this study (eg, hypertension, dyslipide-mia, diabetes, and smoking) are also associated with an increased risk of cancer, Alzheimer disease, lung disease, and renal disease, which combined with CVD deaths accounts for 7 of the 10 leading causes of

death in this age group.⁵³ Finally, all-cause mortality is an objective outcome that is not affected by misclassification bias, which is a significant concern in the determination for cause of death (especially CVD death) among older individuals who typically have multiple chronic diseases.^{54,55}

Limitations of our analysis include that patients in this study may have been healthier than patients referred for a non—exercise stress test, which may limit the generalizability of our results. However, individuals deemed unsuitable for a treadmill stress test are likely to have an even lower fitness level and higher mortality rate. Submaximal cardiorespiratory fitness testing is an

TABLE 2. Hazard Ratios and 95% CIs of All-Cause Mortality by Number of Risk Factors and METs Groups										
No. of risk factor	rs Unadjusted	P value	Model I	P value	Model 2	P value				
0	Reference		Reference		Reference					
I	1.16 (0.98-1.36)	.09	1.14 (0.97-1.35)	.12	1.06 (0.89-1.26)	.53				
2	1.08 (0.92-1.27)	.35	1.12 (0.95-1.31)	.19	1.07 (0.90-1.28)	.46				
≥3	1.10 (0.92-1.30)	.30	1.17 (0.98-1.39)	.08	1.18 (0.97-1.43)	.10				
METs group	Unadjusted	P value	Model I	P value	Model 2	P value				
<6	Reference		Reference		Reference					
6-9.9	0.60 (0.55-0.66)	<.001	0.61 (0.56-0.67)	<.001	0.62 (0.57-0.68)	<.001				
≥ 10	0.43 (0.38-0.49)	<.001	0.43 (0.38-0.49)	<.001	0.44 (0.38-0.50)	<.001				

METs, metabolic equivalents of task.

Model 1: age, sex, race, reason for stress test.

Model 2: model I plus hypertensive medication use, statin use, and diabetes medication use.

alternative measurement of fitness that improves mortality risk stratification and could be considered among individuals unable to perform a traditional treadmill stress test.⁵⁶ In our risk factor-based analyses, we did not examine how well an individual's CVD risk factors were controlled, although adjustment for risk factor treatment with medication did not significantly change the results. In addition, we did not examine change in fitness over time, although we have previously demonstrated that most individuals have a stable relative fitness level during at least 4 years of follow-up.⁵⁷ Strengths of this analysis include that more than 50% of patients were women, and patients were observed for an average of 9.4 years. Survival was determined by Social Security Death Master File, which is nearly identical to the Centers for Disease Control and Prevention methodology for calculating national death statistics in the United States during this time period. In addition, fitness was directly measured using the Bruce protocol, which is the most commonly used method for measuring fitness in clinical practice.

CONCLUSION

Among persons aged 70 years and older who performed a treadmill stress test, an individual's fitness level but not the CVD risk factor burden was significantly associated with long-term mortality risk across the spectrum of CVD risk factor burden. In addition, a high fitness level identified older patients with good long-term survival, even in the presence of a significant CVD risk factor burden. These findings provide added support for the measurement of fitness as a simple and low-cost method to improve risk stratification among older persons.

SUPPLEMENTAL ONLINE MATERIAL

Supplemental material can be found online at http://www.mayoclinicproceedings.org. Supplemental material attached to journal articles has not been edited, and the authors take responsibility for the accuracy of all data. Abbreviations and Acronyms: CAC, coronary artery calcium; CVD, cardiovascular disease; EMR, electronic medical record; HR, hazard ratio; METs, metabolic equivalents of task; PCE, pooled cohort equation

Affiliations (Continued from the first page of this article.): State University, Winston-Salem, NC (P.A.M.); the Department of Medicine, Vanderbilt University Medical Center, Nashville, TN (O.A.O.); the Division of Cardiovascular Medicine, Henry Ford Hospital, Detroit, MI (C.A.B., J.K.E., S.J.K.); and the Houston Methodist DeBakey Heart & Vascular Center, Houston, TX (M.H.A.-M.).

Potential Competing Interests: The authors report no competing interests.

Correspondence: Address to Seamus P. Whelton, MD, MPH, 600 North Wolfe St, Blalock 524A, Baltimore, MD 21287 (seamus.whelton@jhmi.edu; Twitter: @seamuswhelton).

ORCID

Seamus P. Whelton: D https://orcid.org/0000-0001-5914-0735; Paul A. McAuley: D https://orcid.org/0000-0002-9549-7202; Erin D. Michos: D https://orcid.org/0000-0002-5547-5084; Clinton A. Brawner: D https://orcid.org/0000-0002-1705-6620

REFERENCES

- Whelton SP, Roy P, Astor BC, et al. Elevated high-sensitivity Creactive protein as a risk marker of the attenuated relationship between serum cholesterol and cardiovascular events at older age. The ARIC Study. Am J Epidemiol. 2013;178(7):1076-1084.
- Lewington S, Whitlock G, Clarke R, et al. Blood cholesterol and vascular mortality by age, sex, and blood pressure: a metaanalysis of individual data from 61 prospective studies with 55,000 vascular deaths [erratum appears in *Lancet.* 2008; 372(9635):292]. *Lancet.* 2007;370(9602):1829-1839.
- Dalton JE, Rothberg MB, Dawson NV, Krieger NI, Zidar DA, Perzynski AT. Failure of traditional risk factors to adequately predict cardiovascular events in older populations. J Am Geriatr Soc. 2020;68(4):754-761.
- Lloyd-Jones DM, Leip EP, Larson MG, et al. Prediction of lifetime risk for cardiovascular disease by risk factor burden at 50 years of age. *Circulation*. 2006;113(6):791-798.
- Lloyd-Jones DM, Dyer AR, Wang R, Daviglus ML, Greenland P. Risk factor burden in middle age and lifetime risks for cardiovascular and non-cardiovascular death (Chicago Heart Association Detection Project in Industry). Am J Cardiol. 2007;99(4):535-540.
- Benry JD, Dyer A, Cai X, et al. Lifetime risks of cardiovascular disease. N Engl J Med. 2012;366(4):321-329.
- Mortensen MB, Falk E. Primary prevention with statins in the elderly. J Am Coll Cardiol. 2018;71(1):85-94.
- Karmali KN, Goff DC Jr, Ning H, Lloyd-Jones DM. A systematic examination of the 2013 ACC/AHA pooled cohort risk assessment tool for atherosclerotic cardiovascular disease. J Am Coll Cardiol. 2014;64(10):959-968.
- 9. DeFilippis AP, Young R, McEvoy JW, et al. Risk score overestimation: the impact of individual cardiovascular risk factors and preventive therapies on the performance of the American Heart Association—American College of Cardiology—Atherosclerotic Cardiovascular Disease risk score in a modern multi-ethnic cohort. Eur Heart J. 2017;38(8):598-608.
- Murakami Y, Hozawa A, Okamura T, Ueshima H. Evidence for Cardiovascular Prevention From Observational Cohorts in Japan Research Group (EPOCH-JAPAN). Relation of blood

pressure and all-cause mortality in 180,000 Japanese participants: pooled analysis of 13 cohort studies. *Hypertension*. 2008;51(6):1483-1491.

- Hansen TW, Jeppesen J, Rasmussen S, Ibsen H, Torp-Pedersen C. Ambulatory blood pressure and mortality: a population-based study. *Hypertension*. 2005;45(4):499-504.
- Lakka HM, Laaksonen DE, Lakka TA, et al. The metabolic syndrome and total and cardiovascular disease mortality in middleaged men. JAMA. 2002;288(21):2709-2716.
- Doran B, Guo Y, Xu J, et al. Prognostic value of fasting versus nonfasting low-density lipoprotein cholesterol levels on longterm mortality: insight from the National Health and Nutrition Examination Survey III (NHANES-III). *Circulation*. 2014;130(7): 546-553.
- Carter BD, Abnet CC, Feskanich D, et al. Smoking and mortality—beyond established causes. N Engl J Med. 2015;372(7): 631-640.
- Hung RK, Al-Mallah MH, McEvoy JW, et al. Prognostic value of exercise capacity in patients with coronary artery disease: the FIT (Henry Ford Exercise Testing) project. *Mayo Clin Proc.* 2014;89(12):1644-1654.
- Al-Mallah MH, Juraschek SP, Whelton S, et al. Sex differences in cardiorespiratory fitness and all-cause mortality: the Henry Ford Exercise Testing (FIT) Project. *Mayo Clin Proc.* 2016;91(6):755-762.
- Wickramasinghe CD, Ayers CR, Das S, de Lemos JA, Willis BL, Berry JD. Prediction of 30-year risk for cardiovascular mortality by fitness and risk factor levels: the Cooper Center Longitudinal Study. *Circ Cardiovasc Qual Outcomes*. 2014; 7(4):597-602.
- 18. Stamatakis E, Hamer M, O'Donovan G, Batty GD, Kivimaki M. A non-exercise testing method for estimating cardiorespiratory fitness: associations with all-cause and cardiovascular mortality in a pooled analysis of eight population-based cohorts. *Eur Heart J.* 2013;34(10):750-758.
- Kodama S, Saito K, Tanaka S, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. JAMA. 2009;301(19):2024-2035.
- Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. N Engl J Med. 2002;346(11):793-801.
- Ozemek C, Laddu DR, Lavie CJ, et al. An update on the role of cardiorespiratory fitness, structured exercise and lifestyle physical activity in preventing cardiovascular disease and health risk. *Prog Cardiovasc Dis.* 2018;61 (5-6):484-490.
- Bonikowske AR, Lopez-Jimenez F, Barillas-Lara MI, et al. Added value of exercise test findings beyond traditional risk factors for cardiovascular risk stratification. *Int J Cardiol.* 2019;292: 212-217.
- Fleg JL, Morrell CH, Bos AG, et al. Accelerated longitudinal decline of aerobic capacity in healthy older adults. *Circulation*. 2005;112(5):674-682.
- Al-Mallah MH, Keteyian SJ, Brawner CA, Whelton S, Blaha MJ. Rationale and design of the Henry Ford Exercise Testing Project (the FIT project). *Clin Cardiol.* 2014;37(8):456-461.
- Fletcher GF, Ades PA, Kligfield P, et al. Exercise standards for testing and training: a scientific statement from the American Heart Association. *Circulation*. 2013;128(8):873-934.
- 26. Fletcher GF, Balady GJ, Amsterdam EA, et al. Exercise standards for testing and training—a statement for healthcare professionals from the American Heart Association. *Circulation*. 2001;104(14):1694-1740.
- Whelton SP, Dardari Z, Marshall CH, et al. Relation of isolated low high-density lipoprotein cholesterol to mortality and cardiorespiratory fitness (from the Henry Ford Exercise Testing Project [FIT Project]). Am J Cardiol. 2019;123(9):1429-1434.
- Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. J Am Stat Assoc. 1958;53:457-481.

- Cox DR. Regression models and life-tables. J R Stat Soc B. 1972; 34(2):187-220.
- Lavie CJ, Ozemek C, Carbone S, Katzmarzyk PT, Blair SN. Sedentary behavior, exercise, and cardiovascular health. *Girc* Res. 2019;124(5):799-815.
- Fletcher GF, Landolfo C, Niebauer J, Ozemek C, Arena R, Lavie CJ. Promoting physical activity and exercise: JACC Health Promotion Series. J Am Coll Cardiol. 2018;72(14):1622-1639.
- 32. Goff DC Jr, Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines [erratum appears in *Circulation*. 2014;129(25, suppl 2):S74-S75]. *Circulation*. 2014; 129(25, suppl 2):S49-S73.
- 33. Goff DC Jr, Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines [erratum appears in J Am Coll Cardiol. 2014;63(25, pt B):3026]. J Am Coll Cardiol. 2014;63(25, pt B):2935-2959.
- 34. Gransbo K, Almgren P, Nilsson PM, Hedblad B, Engstrom G, Melander O. Risk factor exposure in individuals free from cardiovascular disease differs according to age at first myocardial infarction. *Eur Heart* J. 2016;37(25):1977-1981.
- Sniderman AD, Furberg CD. Age as a modifiable risk factor for cardiovascular disease. Lancet. 2008;371 (9623):1547-1549.
- Wilson PWF, Hoeg JM, D'Agostino RB, et al. Cumulative effects of high cholesterol levels, high blood pressure, and cigarette smoking on carotid stenosis. N Engl J Med. 1997;337(8):516-522.
- Shaw LJ, Raggi P, Berman DS, Callister TQ. Coronary artery calcium as a measure of biologic age. *Atherosclerosis*. 2006;188(1): 112-119.
- 38. Lewington S, Clarke R, Qizilbash N, Peto R, Collins R; Prospective Studies Collaboration. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies [erratum in Lancet. 2003;361(9362):1060]. Lancet. 2002; 360(9349):1903-1913.
- Hamczyk MR, Nevado RM, Barettino A, Fuster V, Andres V. Biological versus chronological aging: JACC Focus Seminar. J Am Coll Cardiol. 2020;75(8):919-930.
- 40. Blaha MJ, Cainzos-Achirica M, Greenland P, et al. Role of coronary artery calcium score of zero and other negative risk markers for cardiovascular disease: the Multi-Ethnic Study of Atherosclerosis (MESA). *Circulation*. 2016;133(9):849-858.
- Kavousi M, Leening MJG, Nanchen D, et al. Comparison of application of the ACC/AHA guidelines, Adult Treatment Panel III guidelines, and European Society of Cardiology guidelines for cardiovascular disease prevention in a European Cohort. JAMA. 2014;311(14):1416-1423.
- Ross R, Blair SN, Arena R, et al. Importance of assessing cardiorespiratory fitness in clinical practice: a case for fitness as a clinical vital sign: a scientific statement from the American Heart Association. *Circulation*. 2016;134(24):e653-e699.
- Clarkson P, Montgomery HE, Mullen MJ, et al. Exercise training enhances endothelial function in young men. J Am Coll Cardiol. 1999;33(5):1379-1385.
- Hambrecht R, Wolf A, Gielen S, et al. Effect of exercise on coronary endothelial function in patients with coronary artery disease. N Engl J Med. 2000;342(7):454-460.
- Kemi OJ, Haram PM, Wisloff U, Ellingsen O. Aerobic fitness is associated with cardiomyocyte contractile capacity and endothelial function in exercise training and detraining. *Circulation*. 2004;109(23):2897-2904.
- 46. Mitnitski AB, Graham JE, Mogilner AJ, Rockwood K. Frailty, fitness and late-life mortality in relation to chronological and biological age. BMC Geriatr. 2002;2:1.
- Koller MT, Leening MJ, Wolbers M, et al. Development and validation of a coronary risk prediction model for older U.

S. and European persons in the Cardiovascular Health Study and the Rotterdam Study. *Ann Intern Med.* 2012;157(6): 389-397.

- Vliegenthart R, Oudkerk M, Hofman A, et al. Coronary calcification improves cardiovascular risk prediction in the elderly. *Circulation*. 2005;112(4):572-577.
- Hippisley-Cox J, Coupland C. Development and validation of QMortality risk prediction algorithm to estimate short term risk of death and assess frailty: cohort study. *BMJ*. 2017;358: j4208.
- Multimorbidity: clinical assessment and management. NICE guideline [NG56]. London: National Institute for Health and Care Excellence; 2016:443.
- Austin PC, van Walraven C, Wodchis WP, Newman A, Anderson GM. Using the Johns Hopkins Aggregated Diagnosis Groups (ADGs) to predict mortality in a general adult population cohort in Ontario, Canada. *Med Care*. 2011;49(10): 932-939.

- van Walraven C. The Hospital-patient One-year Mortality Risk score accurately predicted long-term death risk in hospitalized patients. J Clin Epidemiol. 2014;67(9):1025-1034.
- 53. Heron M. Deaths: leading causes for 2016. Natl Vital Stat Rep. 2018;67(6):1-77.
- Lauer MS, Blackstone EH, Young JB, Topol EJ. Cause of death in clinical research: time for a reassessment? J Am Coll Cardiol. 1999;34(3):618-620.
- Lloyd-Jones DM, Martin DO, Larson MG, Levy D. Accuracy of death certificates for coding coronary heart disease as the cause of death. Ann Intern Med. 1998;129(12):1020-1026.
- de Lannoy L, Sui X, Lavie CJ, Blair SN, Ross R. Change in submaximal cardiorespiratory fitness and all-cause mortality. *Mayo Clin Proc.* 2018;93(2):184-190.
- Brawner CA, Al-Mallah MH, Ehrman JK, Qureshi WT, Blaha MJ, Keteyian SJ. Change in maximal exercise capacity is associated with survival in men and women. *Mayo Clin Proc.* 2017;92(3): 383-390.